

PEST AND DISEASES IN MANGO (*MANGIFERA INDICA* L.)

J. González-Fernández, J.I. Hormaza

IHSM la Mayora CSIC-UMA, 29750 Algarrobo, Malaga, Spain

EXECUTIVE SUMMARY

In this work, we review the most important pests and diseases that affect mango production worldwide as well as the main measures implemented to control them. Pests and diseases are the main factors that can impact sustainable mango fruit production in the tropics and subtropics worldwide. Commercial cultivation of mango, characterized by expansion to new areas, changing crop management, replacement of varieties and increased chemical interventions, has altered significantly the pest and disease community structure in this crop in the different mango producing regions. In addition, climate change is inducing the emergence of new pests and, whereas globalization and trade liberalization have created wide opportunities for mango commercialization growth, at the same time, this can result in faster dispersion of pests and diseases among different mango growing areas if proper sanitary measures are not implemented.

This review covers different topics related to pests and diseases in mango. First, a thorough description of the main pests and diseases that affect mango is provided. Second, the different approaches used in different mango producing countries for chemical and biological control are described. Third, recommendations for appropriate mango management techniques that include integrated pest and disease management, reduction in the use of chemicals and the implementation of a good monitoring and surveillance system to help control the main pests and diseases, are also discussed. Finally, the current knowledge on agrohomeopathy and Korean Natural Farming is analyzed and recommendations on future lines of research to optimize mango pest and disease control are discussed. The fight against mango pests and diseases will require internationally coordinated research, development and innovation efforts to find effective responses and proper management approaches to the extant pests and diseases and be prepared for new threats. This should include the selection of disease and pest tolerant/resistant varieties; the development of those varieties has so far been made through conventional breeding and selection programs and empirical selection made by growers, but new biotechnological approaches will surely speed up this process in the future. Ideally, effective mango pest and disease management will involve a holistic combination of management approaches combined with strict quarantine and regulatory measures that should be enforced for fruit and plant materials at entry points of countries in which mango is produced to prevent introduction of new pests and diseases.

TABLE OF CONTENTS

Introduction	3
Objectives	3
1. Main insect pests affecting mango	4
1.1. Description of the main pests	4
1.1.1. Pests of blossom and tender shoots	11
1.1.2. Pests of fruits	14
1.1.3. Pests of shoots and stems	17
1.1.4. Pests of leaves and buds	18
1.2. Chemical control of the main pests.	19
1.3. Biological control of the main pests.	23
1.4. Mango management approaches to reduce the incidence of pests.	27
2. Main diseases affecting mango	33
2.1. Description of the main diseases	33
2.1.1. Fruit diseases	33
2.1.2. Foliar and floral diseases	35
2.1.3. Soil-borne diseases	39
2.2. Chemical control of the main diseases.	40
2.3. Biological control of the main diseases	44
2.4. Mango management approaches to reduce the incidence of the main diseases.	46
3. Can agrohomeopathy be used to efficiently manage or control pests and diseases in mango?	49
4. What kind of alternative can the KNF (Korean Natural Farming) present as potential strategy to control or manage pests and diseases in mango?	51
5. Future areas and lines of research for the control of mango pests and diseases	52
6. References	54

INTRODUCTION

As in other horticultural crops, pests and diseases are among the main factors that impact sustainable mango fruit production worldwide. Commercial cultivation of mango, characterized by expansion to new areas, changing crop management, replacement of varieties and increased chemical interventions, has altered significantly the pest and disease community structure in this crop. In addition, climate change is resulting in the emergence of new pests and diseases and, whereas globalization and trade liberalization have created wide opportunities for mango commercialization, at the same time, this can result in faster dispersion of pests and diseases among different mango growing areas. As a result, some pests and diseases earlier considered to be minor or secondary have become serious problems recently.

Pests and diseases are serious constraints to mango production throughout the tropics and subtropics. They can affect tree vigor and survival, canopy and root growth, fruit set, yield and pre and post-harvest quality of fruits and have a big impact on the market potential mango fruits. In total, about 400 species of pests are known to infest mango in different parts of the world. Regarding diseases, most plant diseases are caused by fungi. Although anthracnose, malformation and sudden decline are considered as the main mango diseases worldwide, many additional fruit, foliar, floral and soil-borne diseases have been described in different countries and, therefore, could become potential risks in the international mango commercial trade.

Harmonizing efforts at every mango management level as well as combining different approaches are necessary to control and mitigate the destructive effects of pests and diseases on mango production worldwide. Strict quarantine and regulatory measures should be enforced at entry points of countries in which mango is produced to prevent introduction of new pests and diseases. At the farm level, mango growers should be trained to adopt good agricultural practices that include integrated pest and disease management, reduction in the use of chemicals and the implementation of a good monitoring and surveillance system. In addition, coordinated research, development and innovation efforts should be implemented internationally to find effective responses and proper management approaches to the extant pests and diseases and be prepared for new threats.

Although chemical management of mango pests and diseases is widespread, non-chemical measures such as the selection of disease and pest tolerant/resistant varieties are most desirable. The development of those varieties can be made through conventional breeding and selection programs and empirical selection made by growers but also, in recent years, by biotechnological approaches. Ideally, effective mango pest and disease management will involve a holistic combination of management approaches.

In this work we review the most important pests and diseases that affect mango production worldwide as well as the main measures that can be implemented to control them, including biological control. In this sense, a specific section is devoted to evaluating the scarce information available on agrohomeopathy and Korean Natural Farming for mango pest and disease control.

OBJECTIVES

1. To produce an overall picture of the current status worldwide of the main pests and diseases affecting mango (leaves, fruits, inflorescences and branches) with special focus on the countries exporting mango fruits to the US market.
2. To critically review the advances made in biological control of the main pests and diseases affecting mango including agro-homeopathy approaches in order to provide recommendations for biological control management. Special importance will be given to Korean Natural Farming (KNF) approaches that are based on the use of microorganisms to improve the fertility of the soils
3. To define potential future areas or/and lines of research based on the information collected

1. MAIN INSECT PESTS AFFECTING MANGO

1.1. DESCRIPTION OF THE MAIN PESTS

About 400 species of pests are known to infest mango in different parts of the world (Tandon and Verghese 1985; Peña *et al.* 1998). Worldwide lists of pests of mango have been compiled by de Laroussilhe (1980), Tandon and Verghese (1985), Veeresh (1989) and Peña and Mohyuddin (1997). In turn, the pests of mango in India (Srivastava, 1998; Anonymous, 2006), Australia (Anonymous, 1989), Pakistan (Mohyuddin, 1981), Israel (Wysoki *et al.*, 1993; Swirski *et al.*, 2002), the USA (Peña, 1993), Western Africa (Vannière *et al.*, 2004), Brazil (Assis and Rabelo, 2005), Central America (Coto *et al.*, 1995) and Puerto Rico (Martorell, 1975) have also been described.

According to Reddy *et al.* (2018), commercial cultivation of mango, characterized by expansion to new areas, changing crop management, replacement of varieties and increased chemical interventions, has altered the pest community structure significantly. In addition, climate change has induced the emergence of new pests and mango international trade has promoted the movement of pests between regions. As a result, some pests earlier considered to be minor or secondary pests have become serious problems recently (Jayanthi *et al.* 2014a, b).

According to de Faveri (2018), the major pests of mango are mango scale (*Aulacaspis tubercularis*), mango tipborer (*Chlumetia euthysticha*), mango shoot caterpillar (*Penicillaria jocosatrix*), mango cecid flies, MSW (*Sternochetus mangiferae*), mango pulp weevil (*Sternochetus frigidus*), mango red-banded caterpillar (*Deanolis sublimbalis*), mango citripestis (*Citripestis euthraphera*), pink wax scale (*Ceratoplastes rubens*), mango mealybug (*Rastrococcus*), mango planthopper (*Colgaroides acuminata*), mango leafhoppers, tea red spider mite (*Oligonychus coffeae*), tea mosquito bug (*Helopeltis* spp.), mango bud mite (*Aceria mangiferae*), red-banded thrips (*Selenothrips rubricinctus*), flower caterpillars, fruit-spotting bug (*Amblypelta lutescens* and *A. nitida*), coconut bug (*Pseudotheraptus wayi*) and different species of fruit flies (*Bactrocera* spp., *Ceratitis* spp. and *Anastrepha* spp.). Nevertheless, four species on this list, *Chlumetia euthysticha*, *Colgaroides acuminata*, *Amblypelta lutescens* and *A. nitida*, have been so far only reported in Australia.

Below is a summary of arthropod pests of mango, included in Peña and Mohyuddin (1997)

ORDER	FAMILY	SPECIES	GEOGRAPHICAL AREA	PLANT PART AFFECTED
Acari	Eryophidae	<i>Aceria mangiferae</i>	South Pacific, India and Pakistan	Buds
		<i>Cisaberoptus kenyae</i>	Israel	Leaves
		<i>Metaculus mangiferae</i>	Australia and North America	Blossom
	Tarsonemidae	<i>Polyphagotarsonemus latus</i>	North America	Blossom
	Tenuipalpidae	<i>Brevipalpus phoenicis</i>	North America	Leaves
	Tetranychidae	<i>Oligonychus coffeae</i>	Australia	Leaves
		<i>O. mangiferae</i>	India, Pakistan, Israel	Leaves
		<i>O. puniceae</i>	Central America and islands of the Caribbean region	Leaves
		<i>O. yothersi</i>	North America	Leaves
		<i>Tetranychus bimaculatus</i>	Islands of the Caribbean region	Leaves
		<i>T. cinnabarinus</i>	Israel	Leaves
		<i>T. telarius</i>	Southeast Asia	Leaves
		<i>T. tumidus</i>	North America	Leaves
		Tydeidae	<i>Lorrya Formosa</i>	North America
Coleoptera	Bostrichidae	<i>Apate monachus</i>	Islands of the Caribbean region	Trunk and Branches
	Cerambycidae	<i>Batocera rubus</i>	India and Pakistan	Trunk and Branches
		<i>B. rufomaculata</i>	Israel	Trunk and Branches
		<i>Indarbela quadrinonata</i>	India and Pakistan	Trunk and Branches
		<i>Stenodontes downesi</i>	South America	Leaves
		<i>Macrotoma spp.</i>	South America	Trunk and Branches
		<i>M. scutellaris</i>	Israel	Trunk and Branches
	Chrysomelidae	<i>Bassereus brunipes</i>	North America	Leaves
		<i>Crimissa cruralis</i>	South America	Leaves
		<i>Diabrotica balteata</i>	North America	Leaves
		<i>Monolepta lepida</i>	Israel	Leaves
	Curculionidae	<i>Anthonomus spp.</i>	Islands of the Caribbean region	Buds
		<i>Artipus floridanus</i>	North America	Leaves
		<i>Deporaus marginatus</i>	India and Pakistan	Leaves
		<i>Diaprepes abbreviatus</i>	India and Pakistan	Leaves
		<i>Pachneus spp.</i>	North America	Leaves
		<i>Rhynchaenus mangiferae</i>	India and Pakistan	Leaves
		<i>Sternochetus mangiferae</i>	South America, Islands of the Caribbean region	Fruit
	Scarabaeidae	<i>Cotinis nitida</i>	North America	Fruit, Leaves
		<i>Euphoria sepulcralis</i>	North America	Fruit, Leaves
		<i>E. limbata</i>	North America	Fruit, Leaves
		<i>Macraspis spp.</i>		Leaves
		<i>Phyllophaga spp.</i>		Leaves and Root
Scolytidae	<i>Hypocryphalus mangiferae</i>	North and South America	Root, Trunk and Branches	
	<i>Stephanoderes spp.</i>	North America	Trunk and Branches	

Diptera	Cecidomyiidae	<i>Xyleborus saxesini</i>	North America	Fruit
		<i>Procontarina amaramanjae</i>	India and Pakistan	Buds
		<i>P. mangiferae</i>	East Africa and Southeast Asia	Leaves
		<i>P. schreineri</i>	South Pacific	Leaves
		<i>Erosomyia indica</i>	India and Pakistan	Buds and Leaves
		<i>E. mangiferae</i>	Islands of the Caribbean region	Buds
	Loncheidae	<i>Lonchaea spp.</i>	South America	Fruit
	Tephritidae	<i>Anastrepha spp.</i>	South America	Fruit
		<i>A. distincta</i>	South America	Fruit
		<i>A. fraterculus</i>	South America	Fruit
		<i>A. ludens</i>	Central and South America	Fruit
		<i>A. obliqua</i>	Islands of the Caribbean region	Fruit
		<i>A. pseudoparalella</i>	South America	Fruit
		<i>A. serpentina</i>	Central, South and North America	Fruit
		<i>A. striata</i>	Central, South and North America	Fruit
		<i>A. suspensa</i>	Central and North America	Fruit
		<i>Bactrocera jarvisi</i>	Australia	Fruit
		<i>B. aquilonis</i>	Australia	Fruit
		<i>B. carveae</i>	Southeast Asia	Fruit
		<i>B. correcta</i>	Southeast Asia	Fruit
		<i>B. dorsalis</i>	Southeast Asia	Fruit
		<i>B. spp. near B. dorsalis (A)</i>	South Pacific, India and Pakistan	Fruit
		<i>B. spp. near B. dorsalis (B)</i>	Southeast Asia, India and Pakistan	
		<i>B. spp. near B. dorsalis (C)</i>	Southeast Asia, India and Pakistan	
		<i>B. spp. near B. dorsalis (D)</i>	Southeast Asia, India and Pakistan	
		<i>B. facialis</i>	South Pacific	Fruit
		<i>B. frauenfeldi</i>	Australia, Southeast Asia, South Pacific	Fruit
		<i>B. frogatti</i>	South Pacific	Fruit
		<i>B. incisa</i>	Southeast Asia, South Pacific	Fruit
		<i>B. kirki</i>	South Pacific	Fruit
		<i>B. latifrons</i>	Southeast Asia	Fruit
		<i>B. melanota</i>	South Pacific	Fruit
		<i>B. neohumeralis</i>	Australia	Fruit
<i>B. occipitalis</i>		Southeast Asia	Fruit	
<i>B. opilae</i>		Australia	Fruit	
<i>B. passiflorae</i>		South Pacific	Fruit	
<i>B. psidii</i>		South Pacific	Fruit	
<i>B. trilineola</i>	South Pacific	Fruit		
<i>B. tryoni</i>	South Pacific	Fruit		
<i>B. tuberculata</i>	Southeast Asia	Fruit		
<i>B. versicolor</i>	Southeast Asia	Fruit		
<i>B. zonata</i>	South Pacific, Southeast Asia, India and Pakistan	Fruit		
<i>B. (Hemigymnodacus) diversa</i>	Southeast Asia	Fruit		
<i>B. (Zeudacus) cucurbitae</i>	Southeast Asia, India and Pakistan	Fruit		
<i>B. (Zeudacus) tau</i>	Southeast Asia	Fruit		

		<i>B. (Notodacus) xanthodes</i>	South Pacific	Fruit
		<i>Ceratitis capitata</i>	Cosmopolitan	Fruit
		<i>C. catoirii</i>	East Africa	Fruit
		<i>C. cosyra</i>	West Africa	Fruit
		<i>C. punctata</i>	East Africa	Fruit
		<i>C. anonae</i>	East Africa	Fruit
		<i>C. flexuosa</i>	East Africa	Fruit
		<i>C. rosa</i>	East Africa	Fruit
		<i>Dirioxa confuse</i>	Australia	Fruit
		<i>D. pornia</i>	Australia	Fruit
		<i>Cochliomya macellaria</i>	North America	Fruit
		<i>Toxotrypana curvicauda</i>	North, Central and South America, Islands of the Caribbean region	Fruit
Hemiptera	Coreidae	<i>Amblypelta lutescens</i>	Australia	Fruit
		<i>A. nitida</i>	Australia	Fruit
		<i>Pseudotharapterus wayi</i>	East Africa	Fruit
		<i>Veneza stigma</i>	South America	Fruit
	Miridae	<i>Dagbertus fasciatus</i>	North America, Islands of the Caribbean region	Buds
		<i>Rhinacloa spp.</i>	North America. Islands of the Caribbean region	Buds
	Pentatomidae	<i>Brochymena spp.</i>	North America	Leaves
		<i>Plautia affinis</i>	Australia	Leaves
		<i>Stenozygum coloratum</i>	Israel	Fruit?
	Scutelleridae	<i>Symphillus caribbeanus</i>	North America. Islands of the Caribbean region	Fruit
Homoptera	Acanaloniidae	<i>Acanalonia latifrons</i>	North America	Buds, Leaves and Fruit
	Aleyrodidae	<i>Aleurocanthus woglumi</i>	North, Central and South America, West Africa	Leaves
		<i>Aleurodicus dispersus</i>	North America	Leaves
	Aphididae	<i>A. craccivora</i>	Israel	Leaves
		<i>A. fabae</i>	Israel	Leaves
		<i>A. gossypii</i>	Israel	Leaves
		<i>A. spiraeicola</i>	Israel	Leaves
		<i>Toxoptera aurantii</i>	North America, Islands of the Caribbean region	Buds and Leaves
	Asterolecanidae	<i>Asterolecanium pustulans</i>	North America, Israel	Leaves
	Cicadellidae	<i>Amrasca splendens</i>	Australia, East Africa, Southeast Asia, India and Pakistan	Leaves
		<i>Amritodus atkinsoni</i>	India and Pakistan	Leaves
		<i>A. brevistylus</i>	India and Pakistan	Leaves
		<i>Busonomimus manjunathi</i>	India and Pakistan	Leaves
		<i>Chunrocerus niveosparsus</i>	South Pacific	Leaves, Blossom
		<i>Empoasca spp.</i>	Israel	Leaves
		<i>Idioscopus anasuyae</i>	India and Pakistan	Leaves, Fruits
		<i>I. clavosignatus</i>	India and Pakistan	Blossom, Leaves
		<i>I. clypealis</i>	Southeast Asia, India and Pakistan	Leaves, Fruits
		<i>I. decoratus</i>	India and Pakistan	Blossom, Leaves
		<i>I. jayashirae</i>	India and Pakistan	Blossom, Leaves
<i>I. nagpurensis</i>		India and Pakistan	Leaves, Fruits	
<i>I. niveosparsus</i>		Southeast Asia, India and Pakistan	Trunk and Branches,	

			Leaves, Blossom	
		<i>I. nigroclypeatus</i>	India and Pakistan Blossom, Leaves	
		<i>I. spectabilis</i>	India and Pakistan Blossom, Leaves	
		<i>Rabela tabebuia</i>	North America Leaves	
		<i>Scaphytopius spp.</i>	North America Leaves	
	Cixiidae	<i>Myndus crudus</i>	North America Leaves	
	Coccidae	<i>Ceroplastes cirripediformis</i>	North America Leaves	
		<i>C. floridensis</i>	North America, Israel Leaves	
		<i>C. martiniae</i>	South America Trunk and Branches	
		<i>C. rubens</i>	Australia, East Africa, North America, Southeast Asia, Islands of the Caribbean region, India and Pakistan	Leaves
		<i>C. rusci</i>	Israel Leaves	
		<i>C. trochezi</i>	South America Trunk and Branches	
		<i>Coccus acutissimus</i>	North America Leaves	
		<i>C. elatensis</i>	Australia Leaves	
		<i>C. hesperidum</i>	Australia Leaves	
		<i>C. mangiferae</i>	East and West Africa, South and North America, Southeast Asia	Leaves
		<i>C. moestus</i>	Islands of the Caribbean region	Leaves
		<i>C. viridis</i>	North America Leaves	
		<i>Eucalymnatus tessellatus</i>	North America Leaves	
		<i>Kilifa acuminata</i>	North America Leaves	
		<i>Milviscutulus mangiferae</i>	Israel Trunk and Branches, Leaves, Fruits	
		<i>Philephedra tuberculosa</i>	North America Leaves, Fruits	
		<i>Protopulvinaria pyriformis</i>	Israel Leaves	
		<i>P. mangiferae</i>	West Africa Leaves	
		<i>Pulvinaria psidii</i>	Leaves	
		<i>Saissetia oleae</i>	Israel Leaves	
		<i>S. neglecta</i>	Islands of the Caribbean Region Leaves	
		<i>Vinsonia stellifera</i>	South America Leaves	
		Diaspididae	<i>Aonidiella aurantii</i>	Israel Leaves
	<i>A. orientalis</i>		Israel Leaves	
	<i>Aspidiotus destructor</i>		North and South America, Southeast Asia Leaves	
	<i>Aulacaspis tubercularis</i>		Australia, East and West Africa, North and South America, Islands of the Caribbean region	Leaves, Fruits
	<i>Chrysomphalum aonidum</i>		North America, Israel Leaves	
	<i>C. dyctiospermi</i>		North and South America, West Africa, Islands of the Caribbean region	Leaves
	<i>Fiorinia florinae</i>		North America Leaves	
	<i>Hemiberlesia latanidae</i>		North America, Israel Leaves	
	<i>Howardia biclavis</i>		North America Trunk and Branches	
	<i>Ischnaspis longirostris</i>		North and South America Leaves	

		<i>Lindingaspis floridana</i>	North America	Leaves	
		<i>L. ferrisi</i>	India and Pakistan	Leaves	
		<i>Morganella longispina</i>	North America	Trunk and Branches	
		<i>Parlatoria spp.</i>	North America	Leaves	
		<i>Phenacaspis cockerelli</i>	South America	Leaves	
		<i>P. dilatata</i>	Australia	Leaves	
		<i>P. sandwichensis</i>	South Pacific	Leaves	
		<i>Pinnaspis strachani</i>	East Africa, North America	Leaves	
		<i>Pseudaulacaspis cockerelli</i>	North America	Leaves	
		<i>Pseudaonidia trilobitiformia</i>	South and North America	Leaves	
		<i>Radionaspis indica</i>	North America	Leaves	
		<i>Selanaspidus articulatus</i>	South America	Leaves	
		<i>Unaspis citri</i>	South America	Trunk and Branches	
		Flatidae	<i>Colgaroides acuminata</i>	Australia	Blossom, Fruits
		Margarodidae	<i>Drosicha stebbingii</i>	India and Pakistan	Leaves
			<i>D. magiferae</i>	India and Pakistan	Leaves
			<i>Icerya seychellarum</i>	East Africa, Southeast Asia, India and Pakistan	Leaves
		Ortheziidae	<i>Orthezia spp.</i>	South America	Leaves
			<i>O. olivicola</i>	South America	Leaves
		Pseudococcidae	<i>Drosicha mangiferae</i>	India and Pakistan	Leaves
			<i>D. stebbingi</i>	India and Pakistan	Leaves
	<i>Pseudococcus adonidum</i>		East and West Africa, South America, India and Pakistan	Leaves, Fruits	
	<i>P. elisae</i>		South Pacific	Leaves	
	<i>P. longispinus</i>		Islands of the Caribbean region	Buds	
	<i>Rastrococcus invadens</i>		East and West Africa	Buds, Leaves, Fruits	
	<i>R. spinosus</i>		India and Pakistan	Buds, Leaves, Fruits	
	Psyllidae	<i>Apsylla cistellata</i>	India and Pakistan	Leaves	
		<i>Pauropsylla nigra</i>	Southeast Asia	Leaves	
Hymenoptera	Apididae	<i>Trigon spp.</i>	Central America	Blossom	
	Formicidae	<i>Atta spp.</i>	Central America	Leaves	
Isoptera	Termiticidae	<i>Coptotermes acinaciformis</i>	Australia	Root, Trunk and Branches	
		<i>C. formosanus</i>	South Pacific	Trunk and Branches	
		<i>Microcerotermes biroi</i>	Australia	Trunk and Branches	
		<i>M. edenatus</i>		Trunk and Branches	
		<i>Microtermes obesis</i>	India and Pakistan	Trunk and Branches	
		<i>Neotermes insularis</i>	Australia	Trunk and Branches	
		<i>Nisutitermis graveolus</i>	Australia	Root, Trunk and Branches	
		<i>Odontotermes lokanandi</i>	India and Pakistan	Trunk and Branches	
		<i>O. gurdaspurensis</i>	India and Pakistan	Trunk and Branches	

		<i>O. wallonensis</i>	India and Pakistan	Trunk and Branches
		<i>O. obesus</i>	India and Pakistan	Trunk and Branches
		<i>O. horai</i>	India and Pakistan	Trunk and Branches
		<i>Termes cheeii</i>	Australia	Trunk and Branches
Lepidoptera	Arctiidae	<i>Diacrisa obliqua</i>	India and Pakistan	Leaves
		<i>Lymira edwardisii</i>	North America	Leaves
	Coreuthidae	<i>Eccopsis praecedens</i>	East Africa	Fruit
		<i>Lobesia vanillana</i>	East Africa	Fruits
	Ctneuchidae	<i>Syntomeidaepilais jucundisima</i>	North America	Buds
	Gracillariidae	<i>Marmara spp.</i>	North America	Fruits
		<i>Acrocercops spp.</i>	Australia	Leaves
	Gelechiidae	<i>Thiotrina godmani</i>	Islands of the Caribbean region	Buds
	Geometridae	<i>Oxydia spp.</i>	North America	Fruits
		<i>O. vesulia</i>	Islands of the Caribbean region	Fruits
		<i>Pleuroprucha insulsaria</i>	Islands of the Caribbean region	Buds
		<i>Chloropteryx glauciptera</i>	Islands of the Caribbean region	Buds
		<i>Thalassodes dissita</i>	India and Pakistan	Leaves
	Limacodidae	<i>Latoia lepida</i>	Southeast Asia, India and Pakistan	Leaves
	Lymanthriidae	<i>Lymanthria marginata</i>	India and Pakistan	Leaves
	Megatopygidae	<i>Megatopyge defoliata trujillo</i>	North America	Leaves
		<i>M. lanata</i>	South America	Leaves
	Noctuidae	<i>Alabama argillacea</i>	Islands of the Caribbean region	Fruits
		<i>Chlumetia transversa</i>	Southeast Asia, India and Pakistan	Buds
		<i>Gonodonta spp.</i>	Islands of the Caribbean region	Fruits
		<i>G. pyrgo</i>	Islands of the Caribbean region	Fruits
		<i>Othreis fullonia</i>	Australia	Fruits
		<i>O. materna</i>	Islands of the Caribbean region	Fruits
		<i>O. tyrannus</i>	Australia	Fruits
		<i>Penicillaria jocosatrix</i>	South Pacific	Leaves
	Pyalidae	<i>Davara caricae</i>	Islands of the Caribbean region	Buds
		<i>Noorda albizonalis</i>	Southeast Asia, India and Pakistan	Fruits
		<i>Orthaga exvinacea</i>	India and Pakistan	
		<i>Pococera atramentalis</i>	North America, Islands of the Caribbean region	Buds, Fruits
<i>Tallula spp.</i>		North America	Fruits	
Saturnidae	<i>Nataurelia zambesiana</i>	East Africa	Leaves	
Tortricidae	<i>Aethes spp.</i>	Australia, Islands of the Caribbean region	Buds	
	<i>Amorbia aequiflexia</i>	Islands of the Caribbean region	Buds	
	<i>Cosmetra anthophaga</i>	East Africa	Fruits	
	<i>Episimus transferrata</i>	Islands of the Caribbean region	Buds	
	<i>Playnota rostrata</i>	North America	Buds, Fruits	

Orthoptera	Acrididae	<i>Anacridium melanorhodon</i>	East and West Africa	Leaves
Thysanoptera	Paleothripidae	<i>Leothrips sangularis</i>	North America	Buds
	Triptidae	<i>Frankliniella spp.</i>	Islands of the Caribbean region	Buds
		<i>F. bispinosa</i>	North America	Buds
		<i>F. fusca</i>	North America	Buds
		<i>F. kelliae</i>	North America	Buds
		<i>F. occidentalis</i>	Israel	Buds
		<i>Heliothrips hemorroidalis</i>	North America, Israel	Buds
		<i>Scirtothrips mangiferae</i>	Israel	Leaves
		<i>Selenothrips rubrocinctus</i>	North, Central and South America, East Africa, Southeast Asia	Leaves
		<i>Retithrips syriacus</i>	Israel	Leaves
<i>Thrips palmi</i>	North America	Buds		
		<i>T. florum</i>	North America	Buds

Additional insect species considered minor pests in mango in India are included in the following list (Reddy *et al.*, 2018):

Common name	Scientific name	Damage
Blossom feeders and webbers	<i>Asura ruptofascia</i> , <i>Celama analis</i> , <i>C. fasciatus</i> , <i>Cosmostola laesaria</i> , <i>Gymnoscelis imparatalis</i> , <i>Eublemma spp.</i>	Webbing and feeding on the inflorescens
White grub beetles	<i>Holotrichia consanguinea</i> , <i>Anomala sp.</i>	Voracious feeding on leaves during night times
Mango black fly	<i>Aleurocanthus mangiferae</i>	Suck sap from leaves
Painted bug	<i>Coptosoma nazirae</i>	Suck sap from leaves, flowers
Black-eating caterpillar	<i>Indarbela tetraonis</i>	Feed on bark and make holes on stem
Fruit-sucking moths	<i>Eudocima maternal</i> , <i>E. fullonica</i>	Suck sap from fruits

Below we describe with more detail the different pests classified in terms of the tissues affected.

1.1.1. PESTS OF BLOSSOMS AND TENDER SHOOTS

LEAFHOPPERS (Cicadellidae):

Nymphs and adults congregate on panicles and tender shoots where they suck the sap. The continued feeding results in withering and dropping of florets, thus leading to failure of fruit setting. Besides, leafhoppers excrete honey dew which attracts sooty mould and affects photosynthetic efficiency (Butani, 1979). According to Reddy *et al.* (2018), leafhoppers are the major pests of mango with a potential to incur in total fruit losses.

A total of 18 species have been reported as pests affecting mango (Peña *et al.*, 2009); 15 of them have been reported in Asia, but only 3-4 can be considered as relevant (Reddy *et al.*, 2018):

- ***Idioscopus clypealis***: According to Peña (1997), it is present in Southeastern Asia, India and Pakistan; according to EPPO (<https://gd.eppo.int/taxon/IDIOCL/distribution>), it is present in India, Bangladesh and the Philippines, and according to CABI (<https://www.cabi.org/isc/datasheet/28470#toDistributionMaps>), in Australia, Southeastern Asia, India, Pakistan and the South Pacific.

- ***Idioscopus nitidulus*** (*Idiocerus nitidulus*; *Chunrocerus niveosparsus*): it was reported as *Chunrocerus niveosparsus* in Peña (1997) and EPPO, as *I. niveosparsus* in Peña *et al.* (2009) and as *Idioscopus nitidulus* in CABI and Reddy *et al.* (2018). According to Peña (1997) it is distributed in the South Pacific and according to CABI (<https://www.cabi.org/isc/datasheet/28472#toDistributionMaps>) in Australia, Southeastern Asia, India, Pakistan and South Pacific.
- ***Idioscopus nagpurensis***: According to Peña (1997) is present in India and Pakistan and according to CABI also in Southeastern Asia (<https://www.cabi.org/isc/datasheet/28471#toDistributionMaps>)
- ***Amritodus atkinsoni***: reported in Peña (1997) and Reddy *et al.* (2018) in India and Pakistan.

MIDGES (Diptera, Cecidomyiidae)

Mango midges are important pests of mango across the World, but especially in Asia, where about 16 species of gall midges attack mango (Peña, 2002; Ahmed *et al.*, 2005). The main species of this group are:

- ***Erosomya indica*** (mango midge): cited in Peña (1997), Peña *et al.* (2009), Ahmed (2005) and Reddy *et al.* (2018). According to Peña (1997), it is present in India and Pakistan but according to Ahmed (2005) it could be present in other regions as well.
 - ***Erosomya mangiferae*** (mango gall midge or mango blister midge). Cited by Peña (1997), Peña *et al.* (2009) and Reddy *et al.* (2018). According to Peña (1997) it would be present in the Caribbean.
 - ***Asynapta mangiferae***. According to Peña *et al.* (2009) it is present in the Caribbean.
 - ***Gephyraululus mangiferae***. Reported in Peña *et al.* (2009) in the Caribbean.
 - ***Procontarinia matteiana*** (leaf gall midge). According to Peña (1997), it is present in India and Pakistan, and according to Reddy *et al.* (2018), in India, Indonesia, Kenya, Mauritius, Oman, Reunion, South Africa and the United Arab Emirates.
 - ***Procontarinia mangiferae***. According to Peña (1997) it is present in Eastern Africa and Southeastern Asia.
- Procontarinia pustulata***. According to Medina *et al.* (2017) it is present in the Philippines. It has also been reported in northern Australia and Papua New Guinea (Koselik *et al.*, 2009).

THRIPS (Thysanoptera):

Frankliniella occidentalis: reported in Peña (1997), Peña *et al.* (2009) and Reddy *et al.* (2018) in Israel. According to EPPO it would be present in all the continents (<https://gd.eppo.int/taxon/Franoc/distribution>).

Frankliniella bispinosa and ***Frankliniella kellyae***: They have been reported in Peña (1997), Peña *et al.* (2009) and Reddy *et al.* (2018) as mango pests in Florida.

Frankliniella cubensis: reported in Peña *et al.* (2009) in Costa Rica.

Scirtothrips dorsalis: reported in Reddy *et al.* (2018) in India, Thailand and Malaysia. According to EPPO (<https://gd.eppo.int/taxon/SCITDO/distribution>) it is present in India, Pakistan,

Southeastern Asia, Pacific Islands, Australia, some African countries (Ivory Coast, Kenya and Uganda), northern South America and Caribbean.

Thrips palmi: reported by Peña (1997) in North America and by Reddy *et al.* (2018) in India. According to EPPO (<https://gd.eppo.int/taxon/THRIPL/distribution>) it is also present in the Caribbean and several Central and South American countries.

Selenothrips rubrocinctus. Reported by Peña (1997) in North, Central and South America, Eastern Africa and Southeastern Asia. According to CABI (<https://www.cabi.org/ISC/abstract/20056600136>), it is present in Asia (China, Formosa, India, Malaysia, Philippine Islands), Africa (Congo, Fernando Poo, Ghana, Ivory Coast, Nigeria, Principe, Republic of Congo, Sao Thomé, Sierra Leone, Tanganyika, Uganda, Zanzibar), Australasia and Pacific Islands (Hawaii, Mariana Islands, New Caledonia, Papua and New Guinea, Solomon Islands, Wallis Island), Central America (Costa Rica, Honduras, Panama), West Indies and South America (Brazil, British Guiana, Ecuador, Peru, Surinam, Venezuela).

LEPIDOPTERA BLOSSOM FEEDERS:

The lepidopteran flower feeders are considered the second most important inflorescence pests of mango (Peña *et al.*, 2009).

Chloropteryx glauciptera and ***Oxydia vesulia*** (Geometridae). Reported by Peña (1997) and Peña *et al.* (2009) in the Caribbean (Dominica).

Penicillaria jocosatrix (mango shoot caterpillar; Noctuidae). According to Peña (1997) it is present in the South Pacific.

Pococera attramentalis (Pyralidae), ***Pleuroprocha insulsaria*** (Geometridae), ***Platynota rostrana*** (Tortricidae), ***Talulla spp*** (Pyralidae) and ***Racheospila gerularia*** (Geometridae). According to Peña (1997) *P. attramentalis* is present in the Caribbean and *R. gerularia* in Central America. According to Peña *et al.* (2009) some of them could also be present in Florida.

Pleuroprucha asthenaria (Geometridae) and ***Crytoblades gnidiella*** (Pyralidae) reported by Peña *et al.* (2009) in Brazil.

1.1.2. PESTS OF FRUITS:

FRUIT FLIES (Tephitidae):

About 60 different fruit fly species affect mango and related species (Peña *et al.*, 2009). The most important belong to the genera *Anastrepha*, *Bactrocera* and *Ceratitits*.

Anastrepha. The genus is present in the Americas, from the US to Argentina including the Caribbean (Peña, 2009). A total of 12 especies have been reported associated with mango.

- **A. obliqua:** the most common speices in the Americas. According to Peña (1997) and Peña *et al.* (2009), it is present in Australia, Mexico, Costa Rica, Honduras, Guatemala, Cuba, Puerto Rico, Jamaica, El Salvador and Venezuela. According to CABI (<https://www.cabi.org/isc/datasheet/5659#todistributionTable>), in Australia, South America and the islands of the Caribbean.
- **A. striata:** reported by Peña (1997), Peña *et al.* (2009) and CABI (<https://www.cabi.org/isc/datasheet/5667#todistributionTable>) in North, Central, South America and the Caribbean.
- **A. serpentina:** reported by Peña (1997), Peña *et al.* (2009) and CABI (<https://www.cabi.org/isc/datasheet/5665#todistributionTable>) in North, Central, South America and the Caribbean.
- **A. fraterculus:** According to Peña (1997) it is present in South America, according to Peña *et al.* (2009), in Brazil and Ecuador and according to CABI, it is also present in Central America, Mexico and the Caribbean (<https://www.cabi.org/isc/datasheet/5648#todistributionTable>).
- **A. pseudoparalella:** reported by Peña (1997) and Peña *et al.* (2009) in South America.
- **A. suspensa:** reported by Peña (1997) in North and Central America and by CABI (<https://www.cabi.org/isc/datasheet/5668#todistributionTable>) also in the Caribbean and French Guiana in South America.
- **A. ludens:** reported by Peña (1997) in Central and South America and by CABI (<https://www.cabi.org/isc/datasheet/5654#todistributionTable>) in North America (Mexico) and Central America.
- **A. turpiniae** and **A. zuelanie:** reported by Peña *et al.* (2009) in Brazil and Ecuador.

Bactrocera. About 33 species of this genus affect mango in Africa, Asia and Australia. The following are the most important:

- **B. phillipiensis** and **B. occipitalis** according to Peña *et al.* (2009) and CABI (<https://www.cabi.org/isc/abstract/20073277417>) they are present in Asia.
- **B. invadens:** according to Peña *et al.* (2009) present in Western Africa.
- **B. tryoni:** reported by Peña (1997) and Peña *et al.* (2009) in the South Pacific.
- **B. zonata:** reported by Peña (1997) and Peña *et al.* (2009) in Southeast Asia, India, Pakistan and South Pacific and by CABI (<https://www.cabi.org/isc/datasheet/17694#todistributionTable>) in the Near East and Africa.

- ***B. dorsalis***: according to Peña (1997) it is present in the South Pacific and according to CABI (<https://www.cabi.org/isc/datasheet/17685#todistributionTable>) also in the Near East, Eastern Asia, India, Pakistan and Africa.
- ***B. neohumeralis***: reported by Peña (1997) and Peña *et al.* (2009) in Australia and by CABI (<https://www.cabi.org/isc/datasheet/8727#todistributionTable>) in Australia and Papua New Guinea.
- ***B. jarvisi***: reported by CABI (2020) in Australia (<https://www.cabi.org/isc/datasheet/8715>).
- ***B. papayae***: reported by CABI (<https://www.cabi.org/isc/abstract/20063181726>) and The Pacific Community (<https://lrd.spc.int/species/bactrocera-papayae--drew-and-hancock-asian-papaya-fruit-fly>) in Asia (Thailand, Peninsular Malaysia, East Malaysia, Singapore, Indonesia and Kalimantan) and the Pacific (Papua New Guinea).
- ***B. frauenfeldi***: reported by Peña (1997) and Peña *et al.* (2009) in Australia, Southeastern Asia and South Pacific and by CABI (<https://www.cabi.org/isc/datasheet/8712#todistributionTable>) in Australia and South Pacific.

Ceratitis: several species have been reported to affect mango.

- ***C. capitata***: According to CABI (<https://www.cabi.org/isc/datasheet/12367>), it is present in Hawaii, mainland USA, Central and South America, Africa, Spain and other European countries, Australia and the Near East; absent Mexico, the Caribbean and Southeastern Asia.
- ***C. cosyra***: this is one of the three most common species of the genus in Africa. Reported by Peña (1997) and Peña *et al.* (2009) in Western Africa and by CABI (<https://www.cabi.org/isc/datasheet/12370>) in different African countries.
- ***C. fasciventris***: reported in Peña *et al.* (2009) as one of the three most common species of the genus in Africa.
- ***C. anonae***: cited by Peña (1997) and Peña *et al.* (2009) as one of the three most common species of the genus in Africa. According to Peña (1997), present in Eastern Africa.
- ***C. catoirii***: reported by Peña (1997) and Peña *et al.* (2009) in Eastern Africa and Reunion.
- ***C. punctata***: reported by Peña (1997) in Eastern Africa and by CABI (<https://www.cabi.org/isc/datasheet/12376>) in Eastern and Western Africa.
- ***C. flexuosa***: cited by Peña (1997) in Eastern Africa.
- ***C. rosa***: reported by Peña (1997) and CABI (<https://www.cabi.org/isc/datasheet/12378#todistribution>) in Eastern and Southern Africa.
- ***C. silvestrii*** and ***C. quinaria***: reported by Peña *et al.* (2009) in Benin. According to CABI, (<https://www.cabi.org/isc/datasheet/12377>), ***C. quinaria*** would be present in Eastern and Western Africa.

SEED AND PULP WEEVILS:

Mango is not very susceptible to seed and pulp weevils except for some species (Reddy *et al.*, 2018).

Sternochetus mangiferae (seed weevil). Reported by Peña (1997), Peña *et al.* (2009), Reddy *et al.* (2018), EPPO (<https://gd.eppo.int/taxon/CRYPMA>) and CABI (<https://www.cabi.org/isc/datasheet/16434>). It is distributed in most mango growing areas, except North and Central America. A map is available by EPPO (<https://gd.eppo.int/taxon/CRYPMA/distribution>).

Sternochetus frigidus (pulp weevil). According to Peña (1997) and EPPO (<https://gd.eppo.int/taxon/CRYPGR/distribution>), it is present in Southeastern Asia, India, Pakistan and the South Pacific.

***Deanolis sublimbalis* (*Noorda albizonalis*)** (mango seed borer/red-banded Caterpillar/Lepidoptera). According to Peña (1997) and Peña *et al.* (2009), present in Southeastern Asia, India, Pakistan and South Pacific.

Citripestis eutrappera (mango fruit borer (Lepidoptera : Pyralidae). Reported by Reddy *et al.* (2018) in India, Java, Indonesia and Northern Australia.

Deudores isocrates (pomegranate fruit borer/Lepidoptera: Lycaenidae). Reported by Peña *et al.* (2009) in India and the Philippines.

Orgyia postica (cocoa tussock moth/Lepidoptera: Lymantridae). Reported by Peña *et al.* (2009) in the Philippines.

Conogethes punctiferalis (Lepidoptera: Pyralidae): cited by Reddy *et al.* (2018). According to EPPO (<https://gd.eppo.int/taxon/DICHPU/distribution>) it is present in several Asian countries (China, India, Indonesia, Japan, Korea, Malaysia, Myanmar, Sri Lanka, Taiwan and Vietnam), Australia and Papua New Guinea.

OTHER PESTS AFFECTING FRUITS:

Procontarinia frugivora (Diptera, Cecidomyidae). Reported in the Philippines by Gagné & Medina (2004).

Aulacaspis tubercularis (Diaspididae): reported by Peña (1997) and Peña *et al.* (2009) in Australia, Eastern and Western Africa, North and South America and the Caribbean. According to EPPO (<https://gd.eppo.int/taxon/AULSTU/distribution>) it is also present in Europe (Portugal).

1.1.3. PESTS OF SHOOTS AND STEMS

MANGO SHOOT BORERS (Lepidoptera, Noctuidae):

Chlumetia transversa (Lepidoptera: Noctuidae; mango shoot borer). Reported by Peña (1997), in India, Pakistan and Southeastern Asia and according to Reddy *et al.* (2018) in India, Bangladesh, Sri Lanka, Java and the Philippines.

Other shoot borers cited by Reddy *et al.* (2018) include: *Chlumeria alternans* (Noctuidae), *Gatesclarkeana erotias* (Tortricidae; India, Sri Lanka, Timor and Thailand), *Anarsia melanoplecta* (Gelechiidae; India), *A. lineatella* [widespread according to CABI (<https://www.cabi.org/isc/datasheet/5154#todistributionTable>)], *Chelaria spathota* (Gelechiidae) and *Dudua aprobola* [Tortricidae; present in Africa, South and Southeastern Asia and Oceania (http://www.pestnet.org/fact_sheets/mango_flower_webworm_334.htm)].

MANGO STEM BORERS

The mango stem borers are mainly Coleoptera of different families:

Cerambycidae:

- *Batocera rubus* (mango loghorn beetle). Reported by Reddy *et al.* (2018), Peña (1997) and Peña *et al.* (2009) in India and Pakistan and according to EPPO (<https://gd.eppo.int/taxon/BATCRB/distribution>) present in different countries in Asia and the Pacific.
- *B. rufomaculata*. Reported by Peña (1997) in Israel and by Reddy *et al.* (2018) also in India.
- *B. numitor* and *B. titana*. Reported by Reddy *et al.* (2018) at least in India.
- *Glenea multiguttata* and *Coptops aedificator*, reported by Reddy *et al.* (2018) in India.
- *Niphonoclea capito* (mango twig borer). According to Tenorio *et al.* (1989), it is present in the Philippines.

Scolytidae:

- *Hypocryphalus mangiferae* (mango bark beetle). Reported by Peña (1997) and Peña *et al.* (2009) in North and South America, Brazil and Oman and according to Reddy *et al.* (2018), also in Pakistan.
- *Xylosandrus compactus*. Cited by Peña *et al.* (2009) and according to EPPO (https://www.eppo.int/ACTIVITIES/plant_quarantine/alert_list_insects/xylosandrus_compactus), widespread in many countries in America, Europe, Asia and Oceania.

Bostrichidae: *Apate monachus*. Reported by Peña (1997) and Peña *et al.* (2009) in the Caribbean and by CABI (<https://www.cabi.org/isc/datasheet/6071>) also in Africa, India, Brazil and some European countries.

In addition, there are some Homoptera species of the Diaspididae family that can act as stem borers: *Radionaspis indica* and *Morganella longispina*. They were reported by Peña (1997) and Peña *et al.* (2009) in North America. Some reports suggest that the former could be present in Mexico, Panama, Ecuador, the Caribbean, Hawaii, Florida, India, Indonesia and Senegal (<http://scalenet.info/catalogue/Radionaspis%20indica/>) whereas the latter would be widespread (<http://scalenet.info/catalogue/Morganella%20longispina/>).

1.1.4. PESTS OF LEAVES AND BUDS

ARMORED AND SOFT SCALES:

Over 70 scale insects have been reported to affect mango (Reddy *et al.*, 2018).

Aspidiotus destructor (Coccidae). Reported by Reddy *et al.* (2018) in India, Sri Lanka, China, Taiwan, Fiji and Mexico, by Peña (1997) in North and South America and Southeastern Asia and by CABI (<https://www.cabi.org/isc/datasheet/7415>) widespread in different continents.

Drossicha stebbingii (Margarodidae). Reported by Peña (1997) and Peña *et al.* (2009) in India and Pakistan. According to EPPO (<https://gd.eppo.int/taxon/DROCST/distribution>), it is also present in China and Bangladesh.

Rastrococcus invadens (Pseudococcidae). Reported by Peña (1997) and Peña *et al.* (2009) in Eastern and Western Africa and by EPPO (<https://gd.eppo.int/taxon/RASTIN/distribution>) also in Asia.

Aulacaspis tubercularis (Diaspididae): reported by Peña (1997) and Peña *et al.* (2009) in Australia, Eastern and Western Africa, North and South America and the Caribbean. EPPO (<https://gd.eppo.int/taxon/AULSTU/distribution>) increases the number of countries where the pest is present. As described above, it also affects the fruit.

Ceratoplastes rubens (Coccidae). Reported by Peña (1997) in Australia, Eastern Africa, Southeastern Asia, India and Pakistan, North America and the Caribbean. According to EPPO, it is also present in the South Pacific (<https://gd.eppo.int/taxon/CERPRB/distribution>).

There are other scale species such as *Coccus viridis*, *Coccus longulus*, *Ceroplastes actiniformis*, *Philephedra tuberculosa* and the mango shield scales, *Milviscutulus mangiferae* and *Viusonia stellifera*, reported by Peña *et al.* (2009) in Asia, Africa, Australia, Israel and the Americas.

MITES:

Aceria (Eriophyes) mangiferae: cited by Peña (1997), Peña *et al.* (2009) and Reddy *et al.* (2018). It may be a carrier of *Fusarium mangiferae*, which is recognized as the causal agent of mango malformation.

According to this link (http://www.agri.huji.ac.il/mepests/pest/Aceria_mangiferae/) it is present in Egypt and Israel, South Africa, southern Asia, Florida, Mexico and several Central and South American countries.

Oligonychus mangiferae: reported by Peña (1997) in India, Pakistan and Israel and by Venkata (2013) in India and other parts of Asia, Egypt, Mauritius, Peru and Israel.

O. punicae: reported by Peña (1997) in Central America and the Caribbean and by Venkata (2018) also in Australia.

O. coffeae: reported by Reddy *et al.* (2018) in Central America and Australia.

Tetranychus bimaculatus: reported by Peña (1997) in the Caribbean.

Other relevant mite species include ***Metaculus mangiferae***, ***Polyphagus tarsonemus latus***, ***Brevipalpus phoenicis***, ***Oligonychus yothersi*** and ***Tetranychus tumidus*** reported by Peña (1997) in North America.

1.2. CHEMICAL CONTROL OF THE MAIN PESTS

According to Peña *et al.* (2009), the four key pests affecting mango (fruit flies, seed weevils, tree borers and mango leafhoppers) require annual control measures. Secondary pests generally occur at sub-economic levels but can become serious pests as a result of changes in cultural practices and cultivar or because of indiscriminate use of pesticides.

FRUIT FLIES

From the late 1960s, conventional control of fruit flies had involved mainly the use of sprays that combine proteinaceous baits with an insecticide (López *et al.*, 1969; Soto-Manatiú *et al.*, 1987; Mangan *et al.*, 2006; Mangan and Moreno, 2007). For many years the standard insecticide had been malathion (Peck and McQuate, 2000; Burns *et al.*, 2001), although other active ingredients, such as fenthion, deltamethrin, carbaryl or dimethoate, had also been extensively used.

Since the late 1990s, there has been a big effort to find environmentally friendly alternatives to broad-spectrum insecticides (Peck and McQuate, 2000). Growth regulators (such as cyromazine), neonicotinoids (such as imidacloprid), soil microorganisms-derived compounds (such as abamectin and spinosad), plant-derived insecticides (such as azadiractins), and phototoxic dyes (such as Phloxine B) have been successfully tested against various fruit fly species (Díaz-Fleischer *et al.*, 1996; King and Hennessey, 1996; Peck and McQuate, 2000; Vargas *et al.*, 2002; Liburd *et al.*, 2004; McQuate *et al.*, 2005; Díaz-Fleischer *et al.*, 2017). Spinosad-based baits have become one of the most popular alternatives to conventional insecticides in fruit fly control. However, resistance and collateral damage (i.e. negative impact on natural enemies and other beneficial insects or problems with some fruit damage) have been documented (Wang *et al.*, 2005; Hsu and Feng, 2006; Stark *et al.*, 2004; Navarro-Llopis *et al.*, 2012).

Decisions on the time of insecticide applications are usually based on monitoring fruit fly populations with baited traps. In Peru, control measures against *Anastrepha* in mango start when McPhail trap catches average two adults/trap/week (Herrera and Viñas, 1977). In Ecuador, Arias and Jines (2004) recommend a spray of malathion (1%) with protein (4%) once the fruit fly population reaches 0.14 fruit flies/trap/day (FTD). In contrast, studies in Costa Rica, applying at weekly intervals dipterex and malathion reduced damage up to 40% (Soto-Manatiú *et al.*, 1987), while in Mexico, control starts when the fruit is 85-days-old and finishes 2 weeks before harvest (Cabrera *et al.*, 1993).

In spite of the increasing restrictions on their use, insecticides are still widely used among mango growers and are present within many IPM programs for fruit fly control, either sprayed or in alternative strategies such as mass-trapping and attract and kill (AK) techniques, where they are used at a very low volume and inside special devices. This ensures no pesticide application directly onto fruits and lower the risk of environmental contamination. The AK approach is very specific, targeting only the insect pests of interest allowing for highly effective crop protection with just small amounts insecticide that do not get in contact with the edible parts of the plants and minimize the risk to the environment. This method has proven to be an effective and affordable tool to control some fruit fly species, such as *Ceratitis capitata* (Navarro-Llopis *et al.*, 2012; Bouagga *et al.*, 2014).

STONE AND PULP WEEVILS (Curculionidae)

Chemical control has been used with some success for controlling mango weevils in different mango producing countries. In fact, since sanitation practices are usually labor intensive and these pests have few enemies, many mango growers rely mainly on chemical control measures to combat weevil infestations (Chin *et al.*, 2001; Louw, 2009).

When using insecticides, the main approach is to attack diapausing adults by trunk applications or to use foliar sprays at the time of oviposition. During oviposition the adults are active within the canopy, moving onto the fruits and then they can be targeted together with the newly laid eggs. Thus, treatments just before flowering and after initial fruit set have been recommended for seed weevil control (Chin *et al.*, 2010; Prakash, 2012). Several insecticides have been evaluated and recommended (Balock and Kozuma, 1964; Shukla and Tandon, 1985; Louw, 2009; Bajracharya *et al.*, 2012; Reddy *et al.*, 2018). In lab and field tests, effective control of *Sternonchetus mangiferae* was provided by the organochlorine endosulfan, the organophosphates fenthion and malathion, the pyrethroid deltamethrin, the carbamate carbaryl, the phenylpyrazole fipronil and the neonicotinoid thiamethoxam.

Reports on the use of environmentally friendly insecticides for mango weevils control are not abundant and they sometimes show contradictory results. For example, Verghese *et al.* (2004) reported that commercially available azadirachtin was not effective for management of *S. mangiferae* in India, while Bajracharya *et al.* (2012) found that azadirachtin was very effective in reducing pest infestation when compared to control treatment in Nepal.

FRUIT BORERS (Pyralidae)

Control measures against these pests in mango are still mainly depend on the use of pesticides (Istianto and Soemargono, 2015). According to Golez (1991), mango fruit become susceptible to fruit borers 60 days after flowering, and insecticide applications should start at that time. Additional treatments at 75, 90 and 105 days are often required to fully protect the fruit. In Indonesia, Istianto and Soemargono (2015) found that *Noorda albizonalis* began attacking mango fruits when fruits are at the young phase and the attack can occur until the fruits ripened.

Chemicals recommended for control of these pests are deltamethrin and cyfluthrin in the Philippines (Golez, 1991), and fenthion, deltamethrin, indoxacarb and dimethoate in India (Prakash, 2012; Reddy *et al.*, 2018). Plantix recommends sprays with thiacloprid or

chloripyrifos on marble fruit sizes, but always within an integrated approach (<https://plantix.net/en/library/plant-diseases/600128/mango-fruit-borer>). In Indonesia, the application of citronella essential oil reduced the rate of fruit borer attack and the production loss on mango (Istianto and Soemargono, 2015). Repellent, insecticidal, inhibitory, and ovicidal properties of the essential oils of citronella would be behind the effective control of mango fruit borers.

STEM/TRUNK BORERS (Coleoptera)

In India, recommendations for a proper management of mango trunk borers include cleaning the infected holes in the trunks, insert cotton wood soaked in a solution of dichlorvos, close the holes with mud paster, and spray the trunk with chlorpyrifos or imidacloprid or thiamethoxam five times at weekly intervals by changing the chemicals after the onset of monsoon (ICAR, 2014). Also, in India, sprays with carbaryl or quinalphos at fortnight interval from the start of new flushes are recommended (Prakash, 2012). In Nepal, Upadhyay *et al.* (2013) concluded that mango stem borers could be managed by orchard sanitation and destruction of dry shoots from the tree followed by application of imidacloprid or thiomethoxame for 5 times starting from 2nd week of July at 15 days interval. In Florida, USA, pyrethroids have been found to provide control of attacking adults of ambrosia beetles if applied prior to the closing of the galleries with frass (Atkinson *et al.*, 2017).

MANGO LEAFHOPPERS (Hemiptera: Cicadellidae)

Several pesticides have been tried for controlling mango leafhoppers (Tandon and Lai, 1979; Yazdani and Mehto, 1980; Shah *et al.*, 1983; Shukla and Prasad, 1984; Islam and Elegio, 1997; Kudagamage *et al.*, 2001). Khanzada and Naqvi (1985) reported that six sprays of fenitrothion/year were effective in Pakistan. Nachiappan and Baskaran (1986) tested eight insecticides: phasalone, endosulfan, carbaryl, penthoate, fenitrothion, monocrotophos, quinalphos and phosphamidom. Endosulfan provided the best control when spraying was done 1 week after flowering and another treatment 14 days later. Jhala *et al.* (1989) considered that sprays of carbaryl during the off-season maintained the hopper population at low-density levels. Godase *et al.* (2004) demonstrated that sprays of 0.05% monocrotophos at the first panicle emergence and a second spray 15 days later are essential to prevent yield loss. Kudagamage *et al.* (2001) found that imidacloprid controlled mango hoppers if applied just after flowering and again 10 days later. Verghese (2000) recommends using botanical insecticides, like azadirachtin, lemmon grass oil, and citronella oil, if leafhopper populations are low (<4/panicle). If leafhopper density is beyond 4/panicle, he recommends spraying imidacloprid at 0.3 ml/l or thiamethoxam at 0.5 g/l or lambda-cyhalothrin at 0.5 ml/l at panicle initiation stage. In both cases, spraying should be avoided when trees are on full bloom to avoid damage to pollinating insects (Verghese and Devi Thangam 2011).

BLOSSOM / LEAF / TWIG MIDGES (Diptera: Cecidomyiidae)

According to Ahmed *et al.* (2005), midges are commonly controlled by the heavy use of synthetic insecticides, although some less-common techniques to manage populations of these pests have been developed and tested over the last years (Muhammad *et al.*, 2013). In India, spreading chlorpyrifos dust on soil below the tree canopy in April-May, and spraying dimethoate at bud burst stage are recommended (Prakash, 2012; Reddy *et al.*, 2018). In Pakistan, Muhammad *et al.* (2017) found that the insecticides Imidacloprid and Nitenpyram were effective against mango gall midges' larval population, while the trees treated with the insectide Bifenthrin showed least development of galls and concluded that the use of these insecticides can be helpful for controlling mango gall midges.

SOFT AND ARMORED SCALES

In many cases, scale insects become a serious problem in mango orchards following the use of insecticides against other mango pests (Prakash and Patil, 2018). Most scale species can be often suppressed to economic levels by the application of horticultural oils or fish oil resin soap, that dissolves the wax coating and suffocate and kill them (Peña, 2004; Prakash and Patil, 2018). However, this is not always the case, since in India spraying dimethoate at 21 days intervals is a recommendation included in IPM programmes against two mango scales, *Cholopulvinaria polygonata* and *Aspidiotus destructor* (Prakash, 2012). Similarly, Prakash and Patil (2018) found that buprofezin, chlorpyrifos, acephate, lambda-cyhalothrin, profenophos, and dichlorvos were effective insecticides in controlling all stages of the scale *Hemilecanium imbricans* on mango under field conditions, and that the usage of fish oil rosin soap helped in enabling effective penetration of the insecticides.

1.3. BIOLOGICAL CONTROL OF THE MAIN PESTS

FRUIT FLIES

Biological control of fruit flies has mainly relied on parasitoids, especially Braconidae and, at a lesser extent, other families of Hymenoptera (Diapriidae, Chalcididae, Figitidae, Eulophidae, etc.). Classical biological control and augmentative releases of mass-reared parasitoids have been used to suppress *Anastrepha*, *Ceratitis* and *Bactrocera* populations (Wharton, 1978; Sivinski, 1996; Sivinski *et al.*, 1996, 1997, 2000; Montoya *et al.*, 2000). In Florida USA, Mexico, Costa Rica, Brazil, Colombia and Peru, parasitoid species (i.e. *Diachasmimorpha longicaudata* (Ashmead), *Fopius vandenboschi* (Fullaway) and *Aceratoneuromyia indica* (Silvestri)) have been imported and released for the control of *A. suspensa*, *A. ludens* and *A. fraterculus* (Ovruski *et al.*, 2000).

Despite the widespread use of exotic parasitoids over the past 80–100 years, the current trend is to use native species in order to reduce environmental threats (García-Medel *et al.*, 2007; Aluja *et al.*, 2009). This has resulted in numerous studies of the natural enemies of different fruit fly species. For example, Bess *et al.* (1961) reported that the most important parasitoids affecting *Ceratitis capitata* in Hawaii were *Fopius vandenboschi*, *Biosteres oophilus* (= *Opius oophilus*) (= *F. arisanus*) and *B. longicaudatus*. In Brazil, mainly *Doryctobracon areolatus* (Szépligeti) (97%) and *D. longicaudata* (3%) parasitize fruit fly larvae attacking mango (Carvalho and De Queiroz, 2002). In Kenya, Ghana, Tanzania, Uganda and Cote d'Ivoire, the most important parasitoids of *Ceratitis* spp. affecting mango were *Diachasmimorpha fullawayi*, *Fopius caudatus* (Szépligeti), *Psytalia cosyrae* (Wilkinson) and *Tetrastichus giffardianus* Silvestri (Lux *et al.*, 2003). In Mexico and other parts of Latin America, the most common parasitoids attacking fruit flies that affect mangoes (*Anastrepha obliqua*, *A. ludens*, *A. pseudoparallela* and *A. turpiniae*) are *Doryctobracon areolatus*, *Doryctobracon brasiliensis* (Szépligeti), *Doryctobracon crawfordi* (Viereck), *Doryctobracon fluminensis* (Lima) and *Utetes anastrephae* (Viereck) (López *et al.*, 1999; Ovruski *et al.*, 2000; Zucchi, 2000). In Pakistan, the parasitoids attacking *B. zonata* include *Opius longicaudatus* (= *D. longicaudata*), *Dirhinus giffardii* Silvestri, and *Bracon* sp.; *O. longicaudatus* (= *D. longicaudata*), *D. giffardii* and *Spalangia grotiusi* Girault were reported to attack *B. dorsalis*, albeit in small numbers (Syed *et al.*, 1970).

Predators of fruit flies, especially ants, spiders and beetles, have also been identified and some ant species have been used to control fruit flies in mango orchards. Peng and Christian (2006) used the weaver ant, *Oecophylla smaragdina* (Fabricius) for control of *B. jarvisi* in mango orchards in Australia. Van Mele *et al.* (2007) claimed that an African weaver ant (*Oecophylla longinoda*) reduced infestations levels of fruit flies (*Ceratitis* spp. and *Bactrocera dorsalis*) in Benin. The use of these generalist predators could play an important role in biological control of mango pests particularly for resource-poor farmers in developing countries. High populations of natural fruit flies' predators could be linked to a good management of natural weed covers and a proper use of insecticides.

Use of microbial pathogens (fungi, bacteria and nematodes) in fruit fly control has been attempted with varying degrees of success. For example, *Metarhizium anisopliae* has been evaluated in small-scale mango orchards in Kenya and results do not show differences between use of pathogens and use of insecticides (malathion) (Lux *et al.*, 2003). Lezama-Gutierrez *et al.* (2000) also evaluated isolates of *M. anisopliae* against larvae of *A. ludens* with a 22–43%

reduction in adult emergence, depending on the soil where the larvae pupariates. De la Rosa *et al.* (2002) evaluated the fungus *Beauveria bassiana* (Bals.) under laboratory conditions and concluded that the highest control was achieved at the adult stage, while Dimbi *et al.* (2003) reported on the pathogenicity of *M. anisopliae* and *B. bassiana* on different species of *Ceratitis*. Robacker *et al.* (1996) and Toledo *et al.* (1999) tested various strains/isolates of *Bacillus thuringiensis* (Berliner) against larvae of *A. ludens*, *A. obliqua* and *A. serpentina*. Poinar and Hislop (1981), Lindegren and Vail (1986) and Toledo *et al.* (2006) have investigated the use of various nematodes, *Heterorhabditis bacteriophora*, *Heterorhabditis heliothidis* (Khan, Brooks and Hirschmann) and *Steinernema feltiae* Filipjev, against *Anastrepha*, *Bactrocera* and *Ceratitis*. The results were variable for each fruit fly species, with mortalities between 14 and 96%. Some studies suggest that soil type should be considered when selecting the nematode species and planning fruit fly control strategies (Lezama-Gutiérrez *et al.*, 2006).

This biological control of fruit flies can be highly effective and specific when used in combination with autoinoculation devices and insect vectors such as sterile males (FAO/IEA, 2019).

STONE AND PULP WEEVILS (Curculionidae)

Mango weevils have few natural enemies. No parasitoids of *Sternochetus mangiferae* are known (Peña *et al.*, 2009), while some larval parasitoids of *S. frigidus*, viz, *Apanteles* sp., *Angitia trochanterata*, and *Bracon brevicornis*, have been reported (Reddy *et al.*, 2018). Adults may be susceptible to predation by rodents, lizards, birds, and, especially, ants (Hansen, 1993). In fact, in Southastern Asia and Australia, the weaver ant, *Oecophylla smaragdina* (Fabricius), has been reported as an effective predator of some insect pests in mango orchards, including *Sternochetus mangiferae* (Peng and Christian 2007; Van Mele, 2008). In Africa, Abdulla *et al.* (2015) found that *O. longinoda* was effective in suppressing *S. mangifera*.

Regarding control with microorganisms, Shukla *et al.* (1984) reported a baculovirus affecting the larvae of *S. mangiferae*, while in South Africa strains of *Beauveria bassiana* have been tested on mango seed weevil adults and in one laboratory test, two strains caused 30% mortality within 14 days, but in an orchard, neither strain had an effect on the mango seed weevil (Joubert and Labuschagne, 1995).

FRUIT BORERS (Pyralidae)

According to Waterhouse (1998) no natural enemies of mango fruit borers were detected in Java, Indonesia. However, in the Guimaras Islands of the Philippines, the vespid wasp, *Rychium attrisimum*, preys on the larvae, used to stock the wasps' nests as food for their young, as they exit the fruit to pupate and was suspected to contribute to the high larval disappearance in the field. Moreover, the egg parasitoids *Trichogramma chilonis* Ishii and *Trichogramma chilotreae* attack the pest in Luzon (Golez, 1991), and in India, larval parasitoids, such as *Apanteles* sp., *Angitia trochanterata*, and *Bracon brevicornis*, have been reported (Reddy *et al.*, 2018). In spite of this, no references on biological control of these pests through augmentation releases has been found, just the suggestion to maintain natural populations of mango fruit borer predators and parasitoids as high as possible (<https://plantix.net/en/library/plant-diseases/600128/mango-fruit-borer>). This app also recommends, as part of the biological

control approach, apply neem extracts at weekly intervals, starting when mango is flowering and for 2 months.

STEM/TRUNK BORERS (Coleoptera)

Few natural enemies have been reported for suppression of stem and trunk borer populations in mango (Peña *et al.*, 2009). Scheld (1962) recorded a curculionid species (*Scolytopectus schaumii*) which acts as a nest parasite of the ambrosia beetle *Xylosandrus crassiusculus* in the Congo. However, it is unclear whether the ambrosia beetle is killed by the invader, and whether the ambrosia beetle brood continues to develop normally. Most mortality is probably during the dispersal of the adults, and during gallery establishment. The adults of ambrosia beetles are predated by lizards, clerid beetles and ants as they attempt to bore into the host tree (CABI, 2020).

MANGO LEAFHOPPERS (Hemiptera: Cicadellidae)

Several natural enemies of mango leafhoppers have been described from West and Southeastern Asia. Mohyuddin and Mahmood (1993) reported the egg parasitoids, *Gonatocentrus* sp., *Miurfens* sp. nr. *mangiferae* Viggiani and Hayat, *Centrodora* sp. nr. *scolytopae* Valentine, *Aprostocetus* sp. and *Quadrastichus* sp., and the adult ectoparasitoid *Epipyrops fuliginosa* Tames in Pakistan. Fasih and Srivastava (1990) reported that *Aprostocetus* sp., *Gonatocerus* sp. and *Polynema* sp. parasitize eggs. Five species of predators, including *Chrysopa lacciperda* (Kimmins), *Mallada boninensis* (Okomote), *Bochartia* sp. and two unidentified species (one each of Mantidae and Lygaeidae) prey on nymphs (Fasih and Srivastava, 1990). In India, Sadana and Kumari (1991) studied the efficacy of the lyssomanid spider, *Lyssomanes sikkimensis* on *I. clypealis*.

Classical biological control of mango hoppers has not been attempted. Whitwell (1993) described four genera of parasitoids from Dominica, the most common being *Aprostocetus* sp., followed by *Platygaster* sp., *Synopeas* sp. and *Zatropis* sp. Peng and Christian (2005a, b) reported that the weaver ant, *Oecophylla smaragdina* (Hymenoptera: Formicidae) is an efficient biocontrol agent of *I. nididulus* in northern Australia. The entomopathogens, *Verticillium lecanii* (Zimmerman) Viegas, *Beauveria bassiana* Balsamo (Vuillemin) and *Isaria tax*, infect *I. clypealis* in India (Kumar *et al.*, 1993; Srivastava and Tandon, 1986) while the effectiveness of *Metarhizium anisopliae* var. *anisopliae* was tested under laboratory conditions against *A. atkinsoni* (Vyas *et al.*, 1993). Reddy *et al.* (2018) recommends the conservation of natural enemies of mango hoppers, especially coccinellids (e.g. *Coccinella septempunctata*, *C. transversalis* and *Menochilus sexmaculatus*) and spiders. This can be achieved by avoiding spray of broad-spectrum insecticides, and instead entomopathogens like *Metarhizium anisopliae* and botanicals should be used.

BLOSSOM FEEDERS (Lepidoptera)

According to Schreiner (1987), Dipel® reduced caterpillar damage, but careful monitoring or constant spraying was necessary to prevent significant damage. In Brazil, the pesticide *Bacillus*

thuringiensis provided mortality rates of mango caterpillars similar to those of trichlorfon and lambda-cyhalothrin (Barbosa, 2005).

Classical biological control of lepidopteran insects attacking mango in Dominica was initiated with the introduction of the wasps *Aleiodes* sp. and *Euplectrus* sp., and the fly *Blepharella lateralis* Macquart. Populations of the pest were reduced to 25% of pre-release levels; parasitization rates were 20–99%, with *Euplectrus* sp. being the most abundant parasitoid (Nafus, 1991). The parasitoid *Macrocentrus* prob. *delicatus* attacks *Pococera atramentalis*; however, the parasitism rate is unknown (Peña, 1993). In Brazil, *Cryptoblastes gnidiella* is parasitized by *Brachymeria pseudoovata* Blanch (Hymenoptera: Chalcididae), while in Egypt, the endoparasitoid *Tachina larvarum* and the predator *Orius* sp. proved to be good biological control agents against this pest on mango orchards (Kareim *et al.*, 2018).

BLOSSOM / LEAF / TWIG MIDGES (Diptera: Cecidomyiidae)

In India, recommendations for a proper management of cecidomyiid pests in mango include conservation of predators like *Formicai* sp., *Oecophila* sp. and *Camponotus* sp., and parasitoids like *Platygaster* sp., *Systasis* sp. and *Eupelmus* sp., associated with *Dasineura* sp., *Tetrastychus* sp., associated with *E. indica*, and the pteromalid *Pirene* sp., associated with *Procystiphora mangiferae* (Felt) (Reddy *et al.*, 2018). In Pakistan, a survey of midges and their natural enemies associated with mango showed that *Procontarinia* sp. populations were drastically reduced because of increase in parasitism of *Closterocerus pulcherimus* and an unidentified parasitoid (CABI, 2009).

SOFT AND ARMORED SCALES

The populations of most scale species can be often reduced to manageable levels by biological control agents (Peña, 2004; Pradash & Patil, 2018). In fact, conservation of natural enemies (both parasitoids and predators) is a general recommendation for a proper control of scales on mango (Reddy *et al.*, 2018) and, due to the activity of natural enemies, no chemical control or just well-timed insecticide interventions are required (Medina-Urrutia *et al.*, 2017).

A good example of successful biological control of mango scales is *Aulacaspis tubercularis* in South Africa, achieved using the parasitoid *Aphytis chionaspis* and the predator *Cybocephalus binotatus* from Thailand. Close to 50% scale parasitism occurred, while *Cybocephalus binotatus* successfully controlled the remaining scale populations, reducing them to 2%. The release of 500–1000 *Cybocephalus binotatus* beetles per hectare was recommended for effective scale control (Joubert *et al.*, 2000; Le Lagadec, 2004; Daneel and Joubert, 2009). Good control of mango scales have also been achieved using the predator *Aulerodothrips fasciapennis* and the parasitoid *Aspidiotiphagus citrinus* (Kfir and Rosen, 1980, cited by Iyer, 2004). In Ecuador, Arias *et al.* (2004a) observed *Coccidophilus* spp. (Coleoptera: Coccinellidae) and *Chrysopa* spp. preying on *A. tubercularis*; the exotic predator *Cybocephalus nipponicus* (Coleoptera: Nitidulidae) was introduced to supplement predation of the former scale (Arias *et al.*, 2004b).

Mango scale (*Milviscutulus mangiferae*) can be controlled in Israel using parasitoids such as *Coccophagus lycimnia* (Iyer, 2004). The scales *Chrysomphalus aonidum* and *Aonidiella aurantii*

have different wasps as natural enemies: *Aphytis lingnanensis*, *Aphytis holoxanthus*, *Aphytis chrysompali* (Hymenoptera: Aphelinidae) and *Encarsia* spp., which are well distributed in the state of Veracruz, Mexico (Cabrera-Mireles and Ortega-Zaleta, 2004). A wasp, *Anicetus beneficus*, has been identified in Australia for the control of the pink wax scale (*Ceroplastes rubens*) (Cunningham, 1984). Several parasites have been recorded in Israel parasitizing the mango shield scale: *Coccophagus lycimnia* (Walker), *C. eritraensis* Compere, *C. scutellaris* (Dalman), *C. bivittatus* (Compere), *Microterys flavus* (Howard) and *Metaphicus flavus* Howard.

1.4. MANGO MANAGEMENT APPROACHES TO REDUCE THE INCIDENCE OF PESTS

Sampling methods for assessing the status of pest populations has been critical to develop and advance pest control technology (Pedigo and Buntin, 1993). Monitoring and sampling should provide information on pest densities, their dispersion, and dynamics. Through sampling, accurate information is obtained to make accurate decisions which should be based on knowledge of the pest's economic threshold (Peña, 2004). Unfortunately, useful reports on these subjects are scarce or incomplete.

FRUIT FLIES

The combination of several control tools is usually necessary to achieve a good control of fruit flies. A proper orchard management, quarantine treatments or use of the available host resistance could be a good complement to chemical and biological control. A review of some of these approaches is included.

Cultural control

Fruit bagging is reported as one of the best solutions to prevent fruit fly attack of mango and other tropical fruits (Aluja, 1996; Peña *et al.*, 1999). Success with mangoes can be quite high, but more research is needed to determine the type of bags to use for different mango varieties and the best time to bag fruit (Love *et al.*, 2003).

Jirón (1995) reported that *A. obliqua* populations could be reduced by increasing planting distances in order to reduce RH and increase solar radiation within orchards.

A cultural control practice widely recommended and used in many mango producing areas is removal of fallen fruits. In India, this is usually complemented with inter-tree ploughing and raking (followed by insecticide cover sprays), which can reduce fruit fly infestation between 77% and 100% (Verghese *et al.*, 2004).

The use of potassium nitrate (KNO₃) sprays to accelerate and synchronize flowering of mango can, under certain circumstances, help control fruit flies, but it can also exacerbate the problem, so appropriate studies are needed in each producing area.

Mango germplasm

Differences on fruit fly damage between different mango varieties have been found in different fruit fly genus, such as *Bactrocera* and *Anastrepha*. For example, Yee (1987) reported that *B.*

dorsalis does not attack all mango cultivars to the same extent and that the most susceptible cultivars in Hawaii are 'Hawaiian', 'Pirie' and 'Sandersha'. Singh (1991) indicated that the damage caused by *Bactrocera* was highest in fully ripe fruit of 'Mallika' followed by 'Totapari', while frequency of injury in 'Dashehari' ranged from 3.6 to 10% in physiologically mature fruits and ranged from 10 to 25.9% in fully ripe fruit. Susceptibility of different mango cultivars to attack by *A. obliqua* was measured by Carvalho *et al.* (1996) who observed that 'Espada' showed no infestation by *A. obliqua*, whereas 'Carlota' was highly infested. In this study, the survival of adults of *A. obliqua* was lower when the larvae were fed on 'Espada' compared to 'Carlota'. Furthermore, 'Espada' had an adverse effect on the longevity of *A. obliqua* females, possibly due to the presence of toxic substances (Carvalho and De Queiroz, 2002) or absence of essential nutrients. Jirón and Soto-Manitiu (1987) also observed that susceptibility of mangoes to *A. obliqua* differed among cultivars. 'Rosinha', 'Coquinho' and 'Espada' were resistant to *A. obliqua* attack, whereas 'Smith' and 'Pope' were highly susceptible.

According to Joel (1980), the resin ducts in the exocarp of mango fruits confer protection against the vertical movement of the ovipositor and larval movement. Other studies have shown that resistance is related to degree of maturity (Díaz-Fleischer and Aluja, 2003; Aluja and Mangan, 2008); immature mango fruit are less susceptible to *A. suspensa* than mature mangoes when infested artificially (Hennesey and Schnell, 2001). It has been suggested that differences on attack by *A. ludens* to mango might be influenced by volatiles from green or yellow fruits (García-Ramírez *et al.*, 2004).

Quarantine treatments for fruits

Several quarantine treatments have been developed for harvested mangoes. Irradiation, vapor heat or hot water dipping are widely used (Sharp *et al.*, 1988, 1989a, b, c; Hallman and Sharp, 1990; Nascimento *et al.*, 1992; Mangan and Sharp, 1994; Mangan and Hallman, 1998; Shellie and Mangan, 2002a, b; Bustos *et al.*, 2004; additional references in reviews by Mangan and Hallman, 1998 and Follet and Neven, 2006). In fact, in continental USA, quarantine treatments approved for control of fruit flies of the family Tephritidae in mango include irradiation, vapor heat treatments, immersion in hot water, and high temperatures forced air (USDA, 2019). Some of these treatments are only for fruits from some specific countries. For example, vapor heat treatment protocols are developed for fruits from Mexico ("Manila" variety only), the Philippines, and Taiwan. In addition, treatment conditions can be changed according to fruit origin. For instance, minimum irradiation doses are 150 Gy in fruits from Jamaica, Mexico and the Philippines, and 400 Gy in fruits from Dominican Republic, India, Pakistan, Thailand and Vietnam.

Fruit fly monitoring and mass-trapping

Monitoring of fruit flies is crucial to make decisions on when to apply a control or management approach. Moreover, it is key to determine populations dynamics, compare infestation levels and evaluate the effectiveness of a control tactic (Dias *et al.*, 2018). Fruit fly monitoring in mango is mostly performed using traps for adults, because eggs and young larvae are often difficult to see in the fruit and because the primary aim of management programmes is to prevent fruit damage. These traps are also intended for control fruit fly populations when used at high

densities, what is called mass-trapping. Mass trapping networks have contributed to significant reductions in some fruit fly species density (MARNDR, 2014), and have attracted interest due to their efficacy, specificity and low environmental impact.

In the case of *Anastrepha* and some *Bactrocera* species, the most widely used traps since the early 1970s for monitoring and controlling populations are glass and plastic versions of the McPhail trap, which is baited with a mixture of protein and water (Balock and Lopez, 1969; Jirón, 1995). More recently, human urine has been successfully tested as bait for McPhail and McPhail-type traps for resource-poor farmers in tropical countries (Piñero *et al.*, 2003; Aluja and Piñero, 2004). The McPhail trap has provided different results in mango orchards. Balock and Lopez (1969) reported that high concentrations of McPhail traps reduced fly populations and protected mangoes from severe injury during certain periods of the year. However, Aluja *et al.* (1989), working in a mixed mango orchard in Chiapas, Mexico, found that only 31.1% of *Anastrepha* spp. flies landing on the McPhail trap were caught with many flies entering the trap but then escaping. Due to these results and other drawbacks of the McPhail trap, it is being replaced with other types of traps, such as Multi-Lure® traps. Dry synthetic-food-based lures have also been developed, i.e. BioLure® (Suterra LLC, Inc., Bend, Oregon) (Heath *et al.*, 1995, 1997; Epsky *et al.*, 1999) and Nu-Lure® (Advanced Pheromone Technologies) (Robacker and Warfield, 1993; Robacker *et al.*, 1997; Robacker, 2001).

With regard to attractants, methyl eugenol is considered the most powerful male lure for oriental fruit flies. Methyl eugenol was used for successful monitoring, control and eradication of *B. dorsalis* in Oahu Hawaii (Steiner and Lee, 1955), Rota Island (Steiner *et al.*, 1965) and Okinawa, Kume, Miyako and Uaekama Islands, Japan (Iwahashi, 1984). It has been used for monitoring *B. umbrosa* (F.) in the Philippines (Umeya and Hirao, 1975), and is used to lure *B. invadens* in Africa (Lux *et al.*, 2003). In Palau, Pacific Islands, two lures are used to attract mango flies: *Bactrocera fraeunfeldi* (Schiner) is attracted to Cue-lure, and *B. occipitalis* and *B. philippinensis* Drew and Hancock to methyl eugenol (Secretariat of the Pacific Community, 2005). *Bactrocera dorsalis* and *B. umbrosa* were monitored and controlled by mass trapping of males with methyl eugenol and infestations were brought to sub-economic levels in Pakistan (Mohyuddin and Mahmood, 1993). However, concern over the carcinogenicity of methyl eugenol (Waddell *et al.*, 2004) calls for the development of other para-pheromones to attract *Bactrocera* fruit flies. Trimedlure is still considered an important para-pheromone for the Mediterranean fruit fly, with the exception of *C. cosyra* adults, which are attracted to terpinyl acetate and not to trimedlure (Steck, 2003). The attractiveness of mango compounds has been also investigated. For example, some of the volatiles emitted by 'Tommy Atkins' mangoes, i.e. terpenes (*p*-cymene and limonene), are attractive to *C. capitata* adults (Hernández-Sánchez *et al.*, 2001).

The debate is still open with respect to the optimal trap number and distribution and the proper time for trap placement in mango groves. Standardized trapping guidelines for area-wide management of fruit flies recommend 20 to 25 traps per ha for detection surveys (IAEA, 2003). However, Martínez-Ferrer *et al.* (2012) reported that a density of 25 traps per ha was sufficient as a stand-alone method to control the Mediterranean fruit fly (*Ceratitis capitata* Wiedemann; Diptera: Tephritidae) in Spain. Trap densities used in mass-trapping often represent a financial cost that cannot be supported by smallholders in developing countries (Burrack *et al.*, 2008; Lasa *et al.* 2013; Malo & Zapien, 1994), and, therefore, the use of mass-trapping as a control tool

depends on the desing of new cheap attractants and trap devices (Villalobos, 2017). In Haiti, Mertilus *et al.* (2017) evaluated the effectiveness of 2 inexpensive artisanal trap designs as an alternative to the standard McPhail trap for mass trapping fruit flies in mango orchards. They found that the mean number of flies captured in artisanal traps was similar to that captured in the McPhail traps, and that a density of 25 traps per ha was adequate to protect a mango orchard through the maturation phase of the mango season.

Behavioral control

This approach includes two main tactics, Sterile Insect Techique (SIT) and Male Annihilation Technique (MAT).

The SIT involves industrial-scale mass production of radiation-sterilized male insects, which do not lose the ability to fly, mate, and transfer sperm to wild females. Application of fruit fly SIT programmes in mango orchards have been reported in Chile and Brazil for *Ceratitis capitata*, in México for *Anastrepha obliqua* and *A. ludens* (Flores *et al.*, 2014; Flores *et al.*, 2017) and in Thailand for *Bactrocera dorsalis* and *B. correcta* (Sutantawong *et al.*, 2002). SAT still faces challenges, such as the determination of sterile fly release densities requiered to achieve efective sterile wild ratios for the suppression or erraditacion of wild populations (Dias *et al.*, 2018).

MAT involves the distribution of large numbers of dispensers impregnated with a male lure and a toxicant in order to reduce male abundance to such a low level that population suppression or eradication results. Although MAT may be used alone, it is often combined with other control methods, such as the sterile insect releases and/or protein bait sprays. The use of a highly attractive male lure is critical to MAT's effectiveness, and historically it has been most successfully used against ME-responding males and, in particular, the oriental fruit fly, *Bactrocera dorsalis* (Manoukis *et al.*, 2019). The use of MAT against *B. dorsalis* in mango has been reported. In Kenia, Ndlela *et al.* (2016) found that the percentage of infested fruit was 25 and 18 times lower in MAT-treated orchards compared to the control and they recommended that MAT be adopted within a holistic Integrated Pest Management (IPM) approach in the mango agro-system, preferably covering large areas.

STONE AND PULP WEEVILS (Curculionidae)

Field sanitation and quarantine

Orchard hygiene and quarantine treatments are strategies widely recommended and used in mango weevils control programs as a complement to chemical control.

Field sanitation demands complete collection and destruction of fallen fruits and seed material from the orchard, since they will aid in minimising the infestation in following seasons. Trees located near the orchards are often untreated and pose a constant threat of infestation, so it is recommended either removing these trees or using with them a control program to suppress weevils similar to that used in the orchards. Field sanitation is very labour intensive and has been inconsistent in demonstrating efficient pest control (Peña *et al.*, 2009). For instance, in India,

field sanitation reduced infestation of the mango nut weevil, *Sternochetus gravis* (Fabricius), by only 22% (De and Pande, 1987), while in Hawaii, field sanitation failed to reduce infestation rates (Hansen and Armstrong, 1990).

To maintain a mango producing area's weevils free status, common key recommendations are avoid bringing any mango fruit suspected of harbouring weevils within the fruit into the area (for instance, to re-export fruit), and restrictions on movement of plant material from affected sites. Imported mangoes from countries where stone and pulp weevils occur can be subjected to a quarantine treatment. Irradiation is pointed out as the most effective method of killing or sterilizing weevils within fruit (Follet, 2001). In South Africa, irradiation of ripe, marketable fruit protected it from damage and prevented *S. mangiferae* adult emergence (Kok, 1979). In the Philippines, irradiation treatment with a minimum absorbed dose of 165 Gy provides quarantine security for *S. frigidus* in exported mangoes (Obra *et al.*, 2014). In continental USA, minimum irradiation doses approved for control of mango weevils are 165 Gy for *Sternochetus frigidus* in fruits exported from the Philippines, and 300 Gy for *S. mangiferae* in fruits exported from Hawaii, Australia and the Philippines (USDA, 2019). Hot and cold treatment of fruit has also been tried but gave unreliable results and proved phytotoxic (Balock and Kozuma, 1964; Shukla and Tandon, 1985).

FRUIT BORERS (Pyralidae)

General control measures recommended for control of mango fruit borer include collection and destruction of affected and fallen fruits, avoid weed plants that serve as alternative hosts, using light traps (1 per ha) to monitor the activity of adults and fruit bagging at 55-65 days after pollination (Peña *et al.*, 2009; Reddy *et al.*, 2018).

STEM/TRUNK BORERS (Coleoptera)

Orchard sanitation, visual inspections, destruction of dry wood, and pruning and destroying affected branches are general recommendations for the control of this group of pests. In India, stem wrapping with a nylon mesh during May-August helps in capturing freshly emerging adult beetles (Reddy *et al.*, 2014). In addition, a formulation called "sealer cum healer", developed by the Indian Institute of Horticultural Research and applied on the stem along with an insecticide and a fungicide, helps to protect trunks from egg laying by adults (Shivananda *et al.*, 2012). Removing grubs from the infected trunk holes by using iron wire/hook and kill them is also recommended (Reddy *et al.* 2018).

Some success in the detection of ambrosia beetles has been obtained by using traps baited with ethanol in and around port facilities where infested material may be stored, and around nurseries with plants susceptible to attack (CABI, 2019). Nevertheless, the evaluation of new traps as tools for managing ambrosia beetles on mangoes is necessary in order to reduce their damage in newly established groves (Peña *et al.*, 2009).

MANGO HOPPERS (Hemiptera: Cicadellidae)

Reddy *et al.* (2018) recommends avoiding dense planting and maintain tree architecture in such a way that adequate light is penetrated. They also suggest regulating the number of flushes mainly by pruning.

Significant differences in the hopper incidence among genotypes were recorded indicating the scope of host plant resistance (Nachiappan and Bhaskaran 1983; Devi Thangam *et al.*, 2013).

BLOSSOM / LEAF / TWIG MIDGES (Diptera: Cecidomyiidae)

Collection and proper disposal of infested panicles and twigs, and deep ploughing of orchards to expose pupae and diapausing larvae to sun's heat and natural enemies are general recommendations for a proper integrated management of mango cecidomyiid pests in India (Reddy *et al.*, 2018). In Pakistan, Muhammad *et al.* (2013) evaluated the efficiency of colored sticky traps and plastic sheets in capturing adults of *Procontarinia mangicola*. They found that orange colored traps attracted the highest numbers of adults compared to all other traps.

SOFT AND ARMORED SCALES

Pruning to open the tree and a prompt destruction of severely affected leaves and twigs prevent the population buildup of scales on mango (Reddy *et al.*, 2018). These authors also recommend the use of planting material free from scales to minimize the scale population. Removal of attendant ants in order to avoid disturbances on natural enemies of mealy bugs and soft scales is also included in mango IPM programmes (Prakash, 2012).

2. MAIN DISEASES AFFECTING MANGO

Diseases are serious constraints to mango production throughout the subtropics and tropics (Ploetz and Freeman, 2009). They can affect tree vigor and survival, canopy and root growth, fruit set, yield and pre and post-harvest quality of fruit. Although anthracnose, malformation and sudden decline are considered as the main mango diseases worldwide (Ploetz, 2017), many additional fruit, foliar, floral and soil-borne diseases have been described in different countries and, therefore, could become potential risks in the international mango commercial trade. Taking all this into account, we review the most important diseases that affect mango production worldwide, differentiating among those that affect the fruit, the leaves and flowers and soil diseases.

2.1. DESCRIPTION OF THE MAIN DISEASES

2.1.1. FRUIT DISEASES

MAIN FRUIT DISEASES

Among the postharvest diseases of mango fruits, anthracnose is the most prevalent in regions with humid climates, where incidence can affect almost 100% of the fruits. Other common postharvest diseases of mango in humid areas are stem-end rot and black mould rot. Mango black spot is prevalent mainly in dry conditions (Prusky *et al.*, 2009)

Anthracnose

Anthracnose is the most important disease of mango in humid production areas (Lim and Khoo, 1985; Arauz, 2000; Ploetz, 2003, 2018; Ploetz and Freeman, 2009). Although fruit losses can occur in the field, post-harvest losses are most significant (Ploetz, 2018). Anthracnose of mango is caused by species of the genus *Colletotrichum*. In the past, *C. gloeosporioides* (teleomorph: *Glomerella cingulata* (Stoneman) Spauld. & H. Schrenk) was considered the main pathogen associated with this disease (Dodd *et al.*, 1997), but, as the ability to distinguish different species using molecular tools has improved significantly, *Colletotrichum gloeosporioides* has become a species complex with 22 species [e.g. *C. asianum*, *C. fructicola*, *C. gloeosporioides* (*sensu stricto*), *C. queenslandicum*, *C. theobromicola* and *C. tropicale*] (Weir *et al.*, 2012)]. A mango biotype in the *Colletotrichum gloeosporioides* complex, *C. asianum*, may be the most important species (Ploetz, 2018). To date, *C. asianum* has been reported in Australia, Brazil, Florida (USA), Ghana, Mexico, Panama, Philippines, South Africa and Thailand (Honger *et al.*, 2014; Lima *et al.*, 2013; Sharma *et al.*, 2013; Tarnowski, 2009; Udayanga *et al.*, 2013; Weir *et al.*, 2012). Other species complexes, such as *C. acutatum* (*C. fioriniae* and *C. simmondsii*), *C. boninense* (*C. cliviae* and *C. karstii*), *C. siamense* [*C. dianesei* (*syn. C. melanocaulon*) and *C. endomangiferae*], may play secondary roles.

Bacterial black spot (BBS)

In India this disease is known as bacterial canker. This can be the most important mango disease in areas where most other fungi diseases are well managed. Three genetically and pathologically distinct groups of the bacteria responsible of this disease have been identified (Ploetz, 2017a) :

Group I: from Africa, Europe and Asia. *Xanthomonas citri* pv. *mangiferaeindicae* sensu novo.

Group II: from Brazil. *X. axonopodis* pv. *Anacardii*.

Group III. From the French West Indies. *X. axonopodis* pv. *Spondiae*.

Black spot is widespread throughout the world, especially in Asia (Japan, India, Malaysia, Thailand, the Philippines, etc.), Australia, the United Arab Emirates, islands in the Indian Ocean (the Comoros, Reunion, etc.), and East and Southern Africa (Kenya, South Africa, etc.) (https://agritrop.cirad.fr/570027/2/document_570027.pdf.)

Stem-end Rots

The stem-end rot diseases may produce heavy losses during fruit storage and are caused by a diversity of fungal pathogens. According to Prusky *et al.* (2009), *Dothiorella dominicana* (anamorph of *Botryosphaeria dothidea*), *Dothiorella mangiferae*, *Lasiodiplodia theobromae* (*Botryodiplodia theobromae*), *Phomopsis mangiferae* and *Pestalotiopsis mangiferae* would be the dominant pathogens. Ploetz (2018) considered *Lasiodiplodia theobromae* as the most common and widespread pathogen associated with this disease; in addition, *L. pseudoheobromae*, *Neofusicoccum parvum*, *N. mangiferae*, *Botryosphaeria dothidea* and *Neoscytalidium hyalinum* have wide geographical distributions and significantly impact mango.

Alternaria Rot or Black Spot

Alternaria alternata (Fr.) Kreissler (synonyms: *Alternaria fasciculata* (Cooke and Ellis) L. Jones and Grout, *Alternaria tenuis* Nees, and *Macosporiufasciculatum* Cooke and Ellis, no teleomorph known) causes black spot on mango, alternaria leaf spot and lesions on inflorescences (Prusky *et al.*, 1983; Cronje *et al.*, 1990). Although the fungus is cosmopolitan and has a large number of host plants (Neergaard, 1945; Domsch *et al.*, 1980), its effects on mango are most prevalent in arid environments. In Israel, it is a more important disease on fruit than in leaves (Prusky *et al.*, 2009). According to Dodd *et al.* (1997), it has been reported in Australia, Egypt, India, Israel and South Africa.

Black mildew, Sooty Moulds, Sooty Blotch (dothidiomycetes)

Several ascomycetes produce dark, usually superficial growths on stems, leaves and fruits of mango (Ploetz, 2018). Black mildews are caused by a group of tropical obligate plant pathogens (Ploetz, 2018), especially *Meliola mangiferae* (Lim and Khoo, 1985). In contrast, the fungi behind sooty molds are diverse saprophytes that develop in the presence of insects that produce honeydew (e.g., aphids, mealybugs, scales, etc). On the other hand, sooty blotch refers to diseases not associated with honeydew and caused by several groups of dothidiomycetes (Batzer *et al.*, 2005; Johnson *et al.*, 1997; Ploetz *et al.*, 2000).

OCCASSIONAL FRUIT DISEASES

In addition to the main diseases affecting mango fruits, here we include some additional diseases that have been reported occasionally but that rarely cause extensive losses (Snowdon, 1990). Thus, Dodd *et al.* (1997) include: Bacterial rot (*Erwinia* spp.), Blue mold (*Penicillium* spp.), Charcoal rot (*Macrophoma phaseolina* Goidi), Macrophoma rot (*Macrophoma mangiferae* Hingorani and Sharma), Mucor rot (*Mucor* spp.), Phyllosticta rot (*Guignardia mangiferae* Roy.), Phytophthora rot (*Phytophthora nicotianae* var. *parasitica* Waterh.), Rhizopus rot (*Rhizopus* spp.). In addition, Prusky *et al.* (2009) include: Black mold (*Aspergillus* spp.) and Transit rot (*Rhizopus* spp.)

2.1.2. FOLIAR AND FLORAL DISEASES

Algal Leaf Spot (Red Rust)

Algal leaf spot, also known as red rust, is caused by *Cephaleuros virescens* Kunze and, less frequently, by *C. parasiticus* (family Trentepohliaceae, division Chlorophyta (Lim and Khoo, 1985).

Anthracnose (described previously); it was considered previously as a synonym of Blossom Blight but, according to Ploetz (2018) “Blossom blight, which has been attributed to one of the anthracnose agents but is also caused by other fungi”.

Alternaria Leaf Spot (it has been described previously)

Apical necrosis

This disease is caused by a bacterium, *Pseudomonas syringae* pv. *syringae*. It has been reported in Spain, Israel, Portugal and, possibly, Egypt (Cazorla *et al.*, 1998, 2006). Apical buds, leaves and panicles are susceptible, but not the fruit (Cazorla *et al.*, 1998). A dark brown to blackish necrosis develops on vegetative floral buds and can extend from affected buds to leaf petioles causing, in conditions of high damage, the kill of large portions of the canopy and much of the tree's bloom (Ploetz and Freeman, 2009, and Ploetz, 2018)

Bacterial Black Spot (described previously)

Black-banded Disease

This is a relatively unimportant disease that can affect mango leaves and branches (Reddy *et al.*, 1961). The causal fungus, *Rhinocladium corticola* Masee (described as ‘*corticolum*’) (teleomorph: *Peziotrichum corticolum* (Masee) Subramanian), was described on the bark of trees in Poona, India (Hughes, 1980; Prusky *et al.*, 2009).

Black Midew, Sooty Mould and Sooty Blotch (described previously)

Decline Disorders

Diverse biotic and abiotic factors may be primary causes of decline symptoms (McSorley *et al.*, 1980; Kadman and Gazit, 1984; Schaffer *et al.*, 1988; Ploetz and Prakash, 1997). Among the biotic factors, several different fungi cause, or are associated with, decline symptoms worldwide; most are endophytes that have *Botryosphaeria* or *Botryosphaeria*-like teleomorphs (Botryosphaeriaceae). Stress and wound predisposition are usually associated with symptom development (Prusky *et al.*, 2009; Ploetz, 2018). The most important are *Lasiodiplodia theobromae* and *Neofusicoccum parvum*.

The following table from Dodd (1997) shows the main pathogens associated with decline disorders in mango.

LOCATION	KEY REFERENCES	MAJOR CAUSES	PREDISPOSING FACTORS
Australia	Johnson <i>et al.</i> , 1991	<i>Dothiorella dominicana</i>	None
Brazil	Batista, 1947	<i>Diplodia recifensis</i>	Injury, insects
Brazil	Ribiero (1980)	<i>Ceratocystis fimbriata</i>	Injury, insects
Egypt	Acuña and Waite (1977)	<i>Lasiodiplodia theobromae</i>	None
El Salvador	Acuña and Waite (1977)	<i>L. theobromae</i>	Drought, hardpan soil
India	Verma and Singh (1970)	<i>L. theobromae</i>	None
India	Das Gupta and Zacchariah (1945)	<i>L. theobromae</i>	High temperatures
Indonesia	Muller (1940)	<i>L. theobromae</i>	Sun scorch
Malaysia	Lim and Khoo (1985)	<i>L. theobromae</i>	Weakened tres
Niger	Reckhaus and Adamou (1987)	<i>Neoscytalidium dimidiatum</i> = <i>Hendersonula turoloidea</i>	Water stress
Puerto Rico	Alvarez-García et López-García (1987)	<i>L. theobromae</i>	Sun scorch, high RH
South Africa	Darvas (1993)	<i>Dothiorella dominicana</i>	None
USA (Florida)	Ploetz <i>et al.</i> (1996a)	<i>Colletotrichum gloeosporioides</i> , <i>Neofusicoccum parvum</i> , <i>L. theobromae</i> and <i>Phomopsis spp.</i>	Nutritional def. frost, physical damage
USA (Florida)	Ramos <i>et al.</i> (1991)	<i>Botryosphaeria ribis</i>	

Gall and scarly bark

Gall and scarly bark disorders of mango are usually minor problems but can cause a general loss of vigour (Ploetz, 2018). *Fusarium decemcellulare* C. Brick (synonym: *Fusarium rigidiuscula* (Brick) Snyder and Hans.) causes these diseases in Florida (USA), Mexico and Venezuela (Malaguti and de Reyes, 1964; Angulo and Villapudua, 1982; Ploetz *et al.*, 1996b; Prusky *et al.*, 2009).

Grey leafspot

Pestalotiopsis mangiferae (Henn.) Steyaert (synonym: *Pestalolia mangiferae* Henn.; no teleomorph of the fungus is known) causes grey leafspot and stem end rot of mango fruits (Lim and Khoo, 1985; Johnson, 1994b). It is a weak pathogen that usually requires wounding in order to infect mango. Grey leafspot is usually unimportant and occurs mainly on unhealthy or poorly maintained trees. *Pestalotiopsis mangiferae* produces abundant conidia in acervuli that develop in grey leafspot lesions and necrotic areas on fruit (Lim and Khoo, 1985).

Two other species of *Pestalotiopsis* that affect mango are *Pestalotiopsis mangifolia* Guba and *Pestalotiopsis versicolor* Speg. (synonyms: *Pestalotiopsis cliftoniae* Tracy and Earle and *Pestalotiopsis coccolobae* Ellis and Everh.). (Prusky, 2009)

Leaf blight

This disease has been reported in India and Nigeria (Hingorani *et al.*, 1960; Cook, 1975; Okigbo, 2001; Okigbo and Osuinde, 2003), and the causal fungus, *Macrophoma mangiferae* Hingorani and Sharma (Ascomycota), has also been detected in shipments to the USA from Mexico (Systematic Mycology and Microbiology Laboratory, USDA-ARS, Beltsville) although it seems that it is not a serious problem (Prusky, 2009)

Malformation

Malformation is one of the most destructive mango diseases (Ploetz, 2001). Although trees are usually not killed, the vegetative phase of the disease affects canopy development and the infection of the flowers dramatically reduces fruit yield (Prusky *et al.*, 2009).

Malformation was first described in India in 1891 (Kumar and Beniwal, 1991). It is now widely distributed worldwide and continues to spread to the remaining disease-free production areas (e.g. Crespo *et al.*, 2012). To date, the disease has been reported in Australia, Brazil, Myanmar, China, Egypt, El Salvador, India, Israel, Malaysia, Mexico, Nicaragua, Oman, Pakistan, Senegal, South Africa, Sri Lanka, Sudan, Spain, Swaziland, Uganda and the United States (Flechtmann *et al.*, 1973; Crookes and Rijkenberg, 1985; Liew *et al.*, 2016; Lim and Khoo, 1985; Kumar and Beniwal, 1991; Ploetz, 2001; Ploetz and Freeman, 2009; Kvas *et al.*, 2008; Crespo *et al.*, 2012; Senghor *et al.*, 2012; Sinniah *et al.*, 2012; Zhan *et al.*, 2012)

In 1966, *Fusarium moniliforme* was shown to be the agent behind malformation in India (Summanwar *et al.*, 1966). Subsequently, that pathogen, renamed *F. mangiferae* (Britz *et al.*, 2002), has been reported from Australia, China, Egypt, Florida (USA), Israel, Malaysia, Oman, South Africa, Spain and Sri Lanka. Since the description of *F. mangiferae*, a growing list of additional species have also been described as causal agents, including *F. mexicanum* (Otero-Colina *et al.*, 2010) and *F. pseudocircinatum* in Mexico (Freeman *et al.*, 2014a); *F. sterilihyphosum* in Brazil and South Africa (Britz *et al.*, 2002); and *F. tuiense* in Brazil, Senegal and Spain (Freeman *et al.*, 2014b; Lima *et al.*, 2012). In addition, other described [e.g. *F. proliferatum* in Australia, China and Malaysia) and undescribed (in Australia, Mexico and Spain) species in the genus have been associated with the disease (Marasas *et al.*, 2006; Zhan *et al.*, 2010).

Seca and sudden decline

A disease called “Seca” was first reported in 1938 in Pernambuco, Brazil (Viegas, 1960; Rossetto *et al.*, 1996) and later it was also found in the states of Bahia, Goias, the Federal District, Rio de Janeiro and Sao Paulo (Ribeiro, 1997; Colosimo *et al.*, 2000; Silveira *et al.*, 2006). Neighbouring states in Brazil are threatened due to the movement of the pathogen via infected propagation materials and pruning equipment, and a beetle vector (Ploetz, 2018). In 1998, a very similar disease, “sudden decline”, was reported in Oman, Pakistan and the United Arab Emirates (UAE) (Al Adawi *et al.*, 2006). Due to their similarities, they are considered as the same disease in

different mango diseases reviews (Ploetz, 2017). It is a lethal fungal disease. *Ceratocystis fimbriata* Ellis and Halst. *sensu lato* (s.l.) (anamorph: *Thielaviopsis* sp.) was reported in Brazil in the 1930s (Viegas, 1960; Ribiero, 1980; Silveira *et al.*, 2006), and it is recognized as the primary cause of seca. *Diplodia recifensis* Batista (= *Lasiodiplodia theobromae*?) was also reported as the cause of Recife sickness in Brazil (Batista, 1947), but it probably plays no role or a secondary role in the development of this disease (Prusky, 2009).

Two other species have been described for sudden decline in the Oman Gulf Region, *Ceratocystis omanensis*, which is a minor pathogen (Al Subhi *et al.* 2006), and the primary sudden decline agent in Pakistan and Oman, *C. manginecans* (Ploetz, 2018).

Powdery mildew

Powdery mildew is caused by the host-specific fungus *Oidium mangiferae* Berthet (Prakash and Srivistava, 1987; Ploetz and Freeman, 2009). It was first described in Brazil (Berthet, 1914), and it is now reported in most mango producing regions (Palti *et al.*, 1974; Ploetz, 2018).

Phoma blight

Phoma blight caused by *Phoma glomerata* (Corda) Wollenw. and Hochapf (Prakash and Singh, 1977; Prusky *et al.*, 2009) is widespread in India (Prakash and Singh, 1977). It occurs only on old leaves. Initially, lesions are minute and yellow-brown (Prakash and Singh, 1977). As they expand, they darken to brown or cinnamon, become irregular, and may ultimately develop dark margins and dull-grey centres. In severe cases, necrotic patches as large as 13 cm in diameter may develop causing defoliation.

Phoma leafspot

Another *Phoma* sp., *Phoma sorghina* (Sacc.) Boerema. Doren. and Vankest, is the agent behind phoma leafspot, a disease also present in India (Prakash and Singh, 1976). On young leaves, it causes irregular to roughly circular, watersoaked spots, up to 2.5 mm in diameter. Lesions are brown with a yellow to brown margin. Lesions on midribs are elongated and more conspicuous and may coalesce to up to 14 cm in diameter. The symthomps can be confused with those caused by anthracnose (Prusky *et al.*, 2009).

Pink disease

A basidiomycete, *Erythricium salmonicolor* (Berk. and Broome) Burdsall (synonyms: *Corticium salmonicolor* Berk. and Broome, and *Phanerochaete salmonicolor* (Berk. and Broome) Jülich; anamorph: *Necator decretus* Masee) causes pink disease. Pink disease affects many economically important woody plants in the humid tropics, where it is one of the most destructive diseases of mango (Holliday, 1980). The disease is also known as cobweb, rubellosis and thread blight (Prakash and Srivistava, 1987). It has been mainly studied on rubber, *Hevea brasiliensis*, an important host in Southeastern Asia (Rao, 1975). On mango, pink disease can significantly reduce tree vigour and yield, especially in young trees (Lim and Khoo, 1985; Prusky *et al.*, 2009).

Scab

Elsinoë mangiferae Bitancourt and Jenkins (anamorph: *Sphaceloma mangiferae* Bitancourt and Jenkins) causes scab on mango (Bitancourt and Jenkins, 1943; Cook, 1975). The disease was first described in Cuba and Florida (USA) in the 1940s and it is now widespread in the western hemisphere. Scab is important in nurseries, since young tissues are the most susceptible, and because high humidity promotes infection (Ruehle and Ledin, 1955; Prusky *et al.*, 2009).

2.1.3. SOIL-BORNE DISEASES

Soil-borne diseases of mango are relatively less important than foliar and floral diseases, but they can cause significant damage to seedlings, nursery stocks and mature trees.

Phytophthora Diseases

Phytophthora palmivora (E.E. Butler) (Oomycota) causes diseases of mango in several regions of the world: wilt, crown rot, root rot and the death of nursery has been reported in Arizona, the Philippines and Thailand (Kueprakone *et al.*, 1986; Matheron and Matejka, 1988; Tsao *et al.*, 1994); damage to trunks of field-grown, mature trees has been reported in the Ivory Coast (Lourd and Keuli, 1975); damage to fruits in Australia, Malaysia and West Africa (Turner, 1960; Cooke, 2007). Recently, a *Phytophthora* sp. was isolated in Spain from mango trees that were wilted, chlorotic and had sparse canopies and cracked bark (Zea-Bonilla *et al.*, 2007; Prusky *et al.*, 2009).

Root Rot and Damping off

The oomycete *Pythium vexans* de Bary can cause root rot and wilt of seedlings (Lim and Khoo, 1985). Symptoms include wilting of foliage, which initially becomes pale green, but later develops necrotic patches. Roots develop a wet, blackened necrosis that begins in finer roots and progresses to larger roots and the root collar. Prakash and Singh (1980) reported that the basidiomycete *Rhizoctonia solani* Kuhn [teleomorph: *Thanatephorus cucumeris* (Frank) Donk] caused root and damping off of seedlings in India (Prusky *et al.*, 2009)

Sclerotium Rot

The causal fungus is *Sclerotium rolfsii* Sacc. (teleomorph: *Athelia rolfsii* (Curzi) Tu and Kimbrough; synonyms: *Corticium rolfsii* Curzi and *Pellicularia rolftii* E. West) (Prusky *et al.*, 2009). This disease has been reported in Brazil (Almeida *et al.*, 1979), India (Prakash and Singh, 1976) and the Philippines (Palo, 1933).

Verticillium Wilt

The disease was originally attributed to *Verticillium albo-atrum* Reinke and Berth., but later the causing agent became *Verticillium dahliae* Kleb. (Prusky *et al.*, 2009). The disease was first reported in Florida (Marlatt *et al.*, 1970).

White Root Disease

The disease is caused by *Rigidoporus lignosus* (Klotzsch) Imazeki, a basidiomycete that is common in soils of the humid tropics of Africa and Asia (Holliday, 1980). More recently, *Rosellinia necatrix* was reported as the causal agent of White Root Rot in mango trees in Spain (Arjona-Girona and Lopez-Herrera, 2018).

2.2. CHEMICAL CONTROL OF THE MAIN DISEASES

FRUIT DISEASES

ANTHRACNOSE

Although resistance to anthracnose is variable depending on the mango cultivar, even the most tolerant cultivars must be protected by fungicides in humid environments (Lim and Khoo, 1985; Jefferies *et al.*, 1990). In situations where mango fruits develop entirely under disease-favouring conditions, seasonal applications of up to 25 sprays of protective and systemic fungicides have been used (Dodd *et al.*, 1997). However, fungicide use is constrained by the limited number of efficient available products, and by regulations that exist in the producing and/or destination countries (Ploetz, 2018). In general, copper fungicides are the most popular, but their efficacy is often low (Arauz, 2000), and they are usually applied with other fungicides. For example, monthly applications of copper oxychloride combined with mancozeb has been shown as effective for most post-harvest diseases in South Africa (Lonsdale and Kotze, 1993), although the registration of dithiocarbamate fungicides, such as mancozeb, varies among production areas. Preventive treatments with fungicides based on copper or triazoles are the most common decisions of mango growers in Brazil to control anthracnose (Pinto *et al.*, 2004). Another contact fungicide, chlorothalonil, is effective but phytotoxic to fruit larger than a golf ball and, as a result, it should not be used after early fruit set (Ploetz, 2018).

With regard to systemic fungicides, only few are available. The benzimidazoles, primarily benomyl and carbendazim, provided excellent anthracnose control before resistance to them developed (Akem, 2006). Two imidazoles, prochloraz and imazalil, are used in some countries for pre- and post-harvest anthracnose, respectively, since they are moderately effective against this disease, but they are ineffective against stem-end rot (Ploetz, 2018). The strobilurins are effective against anthracnose and several other post-harvest diseases, but to avoid the development of fungicide resistance, no more than three strobilurin applications should be made per season, preferably alternating or combining with fungicides that have a different mode of action (Brent and Hollomon, 2007). Some pre-harvest spray programs used in the control of anthracnose in mango fruits are shown below. Those of Australia, Malaysia and the Philippines are included in Uddin *et al.* (2018), while that proposed for Honduras is described in Huete & Arias (2007).

Country	Pesticides	Number of sprays	Spray timing
Australia	Mancozeb + copper	13	Panicle emergence onwards
Malaysia	Mancozeb + insecticide	Every 10 days	Flower bud onwards
Australia	Prochloraz + copper (applied strategically)	Variable but significantly reduced in dry years	Panicle emergence onwards
Philippines	Mancozeb/chlorothalonil + copper + insecticide	6	Five sprays from induction to fruit set
Honduras	Trifloxystrobin + Propiconazole Thiophanate-methyl Carbendazim Benomyl Mancozeb Chlorotalonil Copper oxychloride	7 Alternating contact and systemics fungicides	Blossom start onwards

Anthracoze forecasting models have been developed to schedule, and reduce, fungicide applications (Fitzell *et al.*, 1984; Dodd *et al.*, 1991). Akem (2006) noted differences between the time prediction of each model; he suggested to use caution when a model was used in an area other than where it was developed. Forecasting would be most useful in seasonally dry situations (where infection occurs only after significant rainfall) (Arauz, 2000). Calendar-based application schedules are needed wherever regular rainfall occurs (Ploetz, 2018).

Fungicide applications usually focus on reducing damage to fruit, but foliar disease control is indicated in some situations and on inflorescences in most situations (Ploetz, 2018). Since infected foliage and branch terminals are important reservoirs of inoculum, fruit set and anthracnose control on fruit are enhanced if applications are made prior to flowering (Jefferies *et al.*, 1990). Off-season control measures are especially beneficial in production environments that receive significant rainfall (Ploetz, 2018). Although pre-harvest sprays and especially fruit-sanitation techniques can eliminate all pathogens on the fruit surface, most of them may have already penetrated the fruit, and, therefore, further treatments to control post-harvest diseases are needed.

Several fungicides have been tested as dip treatments. Benomyl was found effective against quiescent infections of anthracnose of mango in hot water (Peak, 1986), but the application of benomyl after harvest has been banned (Alkan *et al.*, 2018). Post-harvest application of prochloraz in hot and cold dips effectively controls *C. gloeosporioides* and *A. alternata* during storage at low temperature and ripening at 20°C for the cultivars Tommy Atkins, Keitt, Lilly and Haden (Prusky *et al.*, 1999), but it does not provide good control for stem-end-rot (SER). Prochloraz is a well-recognized fungicide that is used commercially to control postharvest

diseases of mango fruit. In Australia, prochloraz at 250 ppm is applied and in Israel it is applied at 300 ppm by overhead spray (Alkan *et al.*, 2018). Other fungicides have been also used successfully for certain mango varieties including thiophanate-methyl and hot imazalil (Secretariat Commonwealth, 1987; Dodd *et al.*, 1991b) The main disadvantage of imidazoles (i.e. prochloraz and imazalil) is that they are less effective at controlling SER pathogens than benzimidazoles (i.e. benomyl and thiabendazole) (Estrada *et al.*, 1996).

With the appearance of various fungicide-resistant isolates, no single fungicide can provide complete protection against anthracnose, alternaria rot and SER, and, consequently, a combination of treatments must be applied to cope with post-harvest pathogens (Alkan *et al.*, 2018). One combination used in Australia is hot water treatment with benomyl followed by a prochloraz spray, which provides effective control of anthracnose, SER and alternaria rot during long storage (Johnson *et al.*, 1990). Another combination applied in Israel includes chlorine sanitation, hot-water brushing (15–20 s) and then a spray of 50-mM hydrochloric acid (HCl), alone or in combination with prochloraz. This combination improved the control of anthracnose and alternaria rot (Prusky *et al.*, 2006). Trials using gamma irradiation to control mango anthracnose have concluded that incorporation of hot fungicide dip is necessary to improve disease control afforded by irradiation [Chadha, 1989]. Appropriate post-harvest treatments have to be selected for individual mango cultivars and possibly even for the same cultivar in different environments (Uddin *et al.*, 2018).

ALTERNARIA ROT

Preharvest treatments with dithiocarbamate fungicides inhibit the development of latent infection. Three sprays with the protectant fungicide maneb, starting 2 weeks after initial fruit set, seem to be most effective (Prusky *et al.*, 1983). However, since quiescent infections do not develop until after harvest and ripening, the application of a postharvest treatment by spraying the fruits on the packing line with prochloraz is simpler and more efficient than the preharvest fungicide treatment (Prusky *et al.*, 2009).

Control of alternaria rot is significantly improved by a combination of physical and chemical treatments. The physical treatment includes a 15–20 seconds hot water spraying and brushing (HWB) treatment at temperatures between 50 and 55°C (Prusky *et al.*, 1999). This approach improved fruit quality and, at the same time, reduced disease incidence. If a prochloraz spray follows this physical treatment it can further improve disease control. Prusky *et al.* (1999) concluded that the type and strength of the postharvest treatment should be optimized according to the level of quiescent infection of *A. alternata* at harvest time. Although prochloraz is very effective for postharvest disease control, a milder postharvest treatment, such as chlorine, can be applied to fruits in which a low incidence of quiescent infections is found at harvest (Prusky *et al.*, 2002). This postharvest physical-chemical treatment has been further improved in light of the finding that *A. alternata* pathogenicity may modulate the pH of the host environment to promote colonization (Eshel *et al.*, 2002; Prusky and Yakoby, 2003; Prusky and Lichter, 2007). Application of a combination of HWB for 15–20 s, followed by spraying with 50 mM hydrochloric acid (HCl), effectively controlled alternaria rot in stored mango fruit. Similar HWB treatments followed by spraying with reduced concentrations of prochloraz at 45 µg/ml in 50 mM HCl inhibited alternaria rot development better than treatment with HCl alone (Prusky *et al.*, 2006). This technology provides a simple treatment for the control of diseases that alkalinize the host environment, including both alternaria rot and anthracnose (Prusky *et al.*, 2009).

STEM END ROT

Postharvest control of *Botryosphaeria* spp. was achieved by postharvest dipping, spraying or ultra-low-volume application of benomyl (where possible). Prochloraz or sodium hypochlorite also effectively suppressed postharvest rot of mango (Plan *et al.*, 2002; Korsten, 2006). A combined treatment of wax and hot water (55°C) provide very effective control of most postharvest pathogens (Sangchote, 1998), but in some cases partial-vacuum infiltration improved disease control, which suggests that control efficiency may have been reduced because the fungicide did not reach the pathogen (Plan *et al.*, 2002).

BACTERIAL BLACK SPOT (BACTERIAL CANKER)

Bacterial Black Spot can be difficult to control on susceptible cultivars, as the available chemicals are marginally effective (Pruvost *et al.*, 1989). During rainy weather, applications of copperbased bactericides are recommended. Their application should focus on protecting fruit and should vary according to the length of the time during which the fruits are exposed to wet conditions (Manicom and Pruvost, 1994). Agricultural antibiotics, such as streptomycin sulphate or nitrate, have been effective (Misra and Prakash, 1992; Viljoen and Kotze, 1972), but their long-term effectiveness is reduced by resistance that develops after continued use.

FOLIAR AND FLORAL DISEASES

SECA AND SUDDEN DECLINE

According to Ploetz (2018), managing these diseases with fungicides is still a challenge, especially on susceptible cultivars. External applications of protectant or systemic fungicides would probably be ineffective given the internal location of the pathogens. In areas where partially resistant cultivars are grown, the removal and burning of affected branches and treatment of the exposed branch stubs with copper fungicides are recommended (Ribeiro *et al.*, 1995; Ribeiro, 1997). Alternatively, injecting fungicides might be effective, as it has been done to control Dutch elm disease (Ploetz, 2018).

MALFORMATION

The internal location of the pathogen in affected trees makes it difficult to control this disease with chemicals (Ploetz, 2018). Thus, although a diverse array of pesticides, hormones and growth regulators have been tested against malformation, only few have shown some potential. In India, spray with carbendazim every fifteen days has been recommended (Misra *et al.*, 2000). In Pakistan, Iqbal *et al.* (2011) found that clipping malformed branches at 45 cm distance followed by spray of benomyl showed the best results with a 70% decrease over previous years damage. Freeman *et al.* (2014b) described the use of prochloraz applications in conjunction with sanitation measures. Although significant reductions in malformation were reported, prochloraz is not approved for use in mango in some producing (i.e. the United States) and importing countries. Darvas (1987) reduced the percentage of malformed inflorescences from 96% to 48% by injecting 'Keitt' trees with the fungicide fosetyl-Al, although no significant increase in fruit yield was observed. Recently, more extensive applications of phosphonates (the active ingredient of fosetyl-Al) were shown to be effective against malformation in South Africa. Nonetheless, the efficacy of intensive phosphonate applications needs to be demonstrated in other areas, and its adoption would need to address phosphonate residue tolerance limits that are imposed in some importing countries (Ploetz, 2018). In Egypt, foliar application of nano-chitosan on mango trees improved the vegetative growth and fruit quality, increased yield and decreased the incidence of malformation (Zagzog *et al.*, 2017).

POWDERY MILDEW

A large number of fungicides have been used against this disease in different mango growing countries. Apart from dormant sprays, several applications of suitable fungicides at 15-20 day-intervals are required to effectively control the disease. Initially, inorganic copper or sulfur-based chemicals were used and then a broad range of organic and systemic fungicides, which acted as eradicants, protectants or both, were introduced (Nasir *et al.* 2014). These authors reviewed the main groups of chemicals used against the powdery mildew of mango: copper-based fungicides, sulphur fungicides, chlorothalonil, nitro compounds, and systemic fungicides (benzimidazoles, imidazole, morpholines, organophosphorus, oxathiins, piperazine, pyridimines, strobilurins and triazoles).

Sulfur fungicides, as dusts or sprays, are widely used and provide reasonable protectant control of powdery mildews (Palti *et al.*, 1974; Gupta and Yadav, 1984; Prakash and Misra, 1986; Kawate, 1993; Prakash and Raof, 1994; Desai, 1998; Chavan *et al.*, 2009), although they can burn flowers and young fruits during warm, sunny conditions (Johnson, 1994a). Systemic fungicides in general are very effective in reducing the disease (Ihsan *et al.*, 1999). In addition, some fungicides, such as dinocap, fenbuconazole and hexaconazole, can reduce pollen germination (Dag *et al.*, 2001), and, consequently, their use should be limited during the flowering season. Application of phosphate solutions is a new and safer approach in the control strategies of powdery mildews in several vegetables and fruit trees (Nasir *et al.*, 2014). In mango, foliar sprays of K₂HPO₄ and KH₂PO₄, especially in alternation with systemic fungicides, were effective against powdery mildew (Nofal and Haggag, 2006; Reuveni *et al.*, 1998).

2.3. BIOLOGICAL CONTROL OF THE MAIN DISEASES

FRUIT DISEASES

ANTHRACNOSE AND STEM-END-ROT

Bacillus licheniformis, on its own or alternated with copper oxychloride (allowed in organic farming management), has been evaluated as a preharvest spray treatment to control mango fruit diseases (Prusky *et al.*, 2009). Preharvest applications of *B. licheniformis* at 3-week intervals from flowering until harvest controlled moderate levels of anthracnose and of soft rot caused by *Botryosphaeria*, which suggests a potential treatment for commercial preharvest applications (Silimela and Korsten, 2007).

Post-harvest biological control agents have been the focus of considerable research (Droby *et al.* 2016). A number of microorganisms with *in vitro* or *in vivo* activity against *C. gloeosporioides* have been isolated (Jeffries and Koomen, 1992), but few examples have been used commercially in the field until Korsten (2004) isolated *Bacillus licheniformis* from leaf and fruit surfaces, and effectively controlled anthracnose of mango. This product was used either alone or in combination with hot water treatments for 5 minutes at 45°C and with low doses of prochloraz or sodium hypochlorite, although only when used alone could be considered in organic agriculture (Govender *et al.*, 2005). The yeasts *Rhodotorula minuta* (Patino-Vera *et al.*, 2005) and *Debaryomyces nepalensis* (Luo *et al.*, 2015) have also been suggested as potential biocontrol agents of anthracnose, but they have not been widely applied commercially (Droby *et al.*, 2016). Other approaches to anthracnose control using biological methods included the use of a non-pathogenic strain of *Colletotrichum magna* that colonizes the fruit endophytically and prevents infection by *C. gloeosporioides* (Prusky *et al.*, 1993), and the expression of an antifungal peptide

in the yeast *Saccharomyces*, which controlled postharvest diseases caused by *C. coccodes* (Jones and Prusky, 2001). Recently, Luo *et al.* (2015) found that the yeast *Debaryomyces nepalensis* decreased the decay incidence to anthracnose while maintained storage quality of mango fruits. In Thailand, Rungjindamai (2016) found that two isolates of epiphytic bacteria, identified as *Bacillus* sp. MB61 and *Bacillus* sp. LB72, reduced the size of the lesions caused by *C. gloeosporioides*.

BACTERIAL BLACK SPOT (BACTERIAL CANKER)

Biological control measures against Bacterial Black Spot have not been widely studied (Prusky *et al.*, 2009). In India, Kishun (1994) indicated that a strain of *Bacillus coagulans* from the phylloplane of mango was effective against strains of the pathogen, although control of bacterial black spot in the field was not reported.

MALFORMATION

Some attempts to control mango malformation through biopesticides have been reported in India. Three different species of *Trichoderma* i.e., *Trichoderma viride*, *T. virens* and *T. harzianum*, were tested against the *F. moniliforme* var. *subglutinans* (Kumar *et al.*, 2012). The three bioagents varied in their efficacy against *F. moniliforme* var. *subglutinans*, but, in general, all of them were effective in decreasing the growth of all evaluated isolates of *Fusarium*. Kumar *et al.* (2009) evaluated antifungal activity against *F. moniliforme* var. *subglutinans* of leaf extracts from different plants. Although all the leaf extracts limited the radial growth of the fungus, extracts of *Azadirachta indica*, *Achyrenthes roseus* and *Calotropis gigantea* were found more effective. Usha *et al.* (2009) also reported antifungal activity of *Datura stramonium*, *Calotropis gigantea* and *Azadirachta indica* against floral malformation pathogens.

In South Africa, several bacterial isolates obtained from mango orchards as well as other environments, were screened for their antifungal properties against *Fusarium* and five (identified as *Alcagenes faecalis*) were able to significantly inhibit the growth of the pathogen (Veldman *et al.*, 2017). The authors suggest that the modes of action of these bacteria involved a combination of competition for space and production of secondary metabolites, such as volatiles, phenolic compounds and siderophores.

POWDERY MILDEW

Sztejnberg *et al.* (1989) reported that an isolate of *Ampelomyces quisqualis* parasitized powdery mildew of mango and reduced the disease in field trials. He also found that *A. quisqualis* was tolerant to many fungicides currently used to control powdery mildew. Nofal and Haggag (2006) reported that in vitro application of biocontrol agents as *Verticillium lecanii*, *Bacillus subtilis* and *Tilletiopsis minor* leaf disks before inoculation with *O. mangiferae* markedly decreased conidial germination and leaf infection. In field trials, the application of those agents at 15 days intervals effectively controlled *O. mangiferae* on blossom clusters and fruit set on naturally powdery mildew infected cultivars Alphonso and Seddek. Mixing kaolin and monopotassium phosphate with biocontrol agents increased their efficacy. In Egypt, Azmy (2014) found that spraying with the bio-fungicide AQ10 (*Ampelomyces quisqualis*) at the rate of 0.005% after harvesting the crop showed good reduction of powdery mildew severity on mango trees with an increase of fruit yield. Kaur *et al.* (2018) found that among six biocontrol agents evaluated by giving three sprays (starting from two weeks after the panicle emergence) at 15 days intervals against powdery in mango, two antagonists, namely *Bacillus subtilis* and *Ampelomyces quisqualis*, exhibited high degree of disease control, when tested over two different locations. The mechanisms implicated

in biological control of powdery mildew fungus include mycoparasitism, antibiosis, competition, and induced resistance.

2.4. MANGO MANAGEMENT APPROACHES TO REDUCE THE INCIDENCE OF DISEASES

FRUIT DISEASES

ANTHRACNOSE, STEM-END ROT AND ALTERNARIA ROT

Pre-harvest control measures

Post-harvest diseases can be reduced by various pre-harvest control measures, including the use of tolerant cultivars, orchard hygiene, manipulation of flowering and integrated management using chemical, physical and biological controls (Johnson *et al.*, 1989; Ploetz, 2004; Akem, 2006).

Since the development of mango anthracnose is dependent on high humidity, mango orchards should ideally be established in areas with a well-defined dry season, to allow for fruit development under conditions unfavourable for disease development (Prusky *et al.*, 2009). In the tropics, mango flowering usually occurs during the dry seasons, and the incidence and severity of mango anthracnose can be close to zero in fruits that develop completely in the dry season, without the need of any additional control measures (Arauz, 2000). However, anthracnose incidence of > 90% is common in fruits that develop during the rainy season (Arauz, 1999). Thus, modifying flowering time to a less sensitive period could be an appropriate option. Flowering can be advanced by several weeks by applying potassium nitrate sprays to mature foliage (Núñez-Elisea, 1985). The growth retardant paclobutrazol, alone or followed by potassium nitrate sprays, can also be used to advance flowering (Núñez-Elisea *et al.*, 1993) although its use is not allowed in some countries.

Sanitation of the tree in the field is a difficult practice since elimination of dry panicles and mummified fruits is time consuming. Bagging can result in reduced anthracnose severity, but it also reduces the red colour of the fruit of some varieties, which could reduce consumer appeal in some markets (Hofman *et al.*, 1997).

Although all commercial mango cultivars are susceptible to anthracnose, some varieties are less susceptible than others. Thus, 'Tommy Atkins' and 'Keitt' seem to be less susceptible than 'Irwin', 'Kent', 'Haden' and 'Edward' (Campbell, 1992). Consequently, cultivar selection should be taken into account in areas with high incidence of the disease.

Regarding stem-end rot, Johnson *et al.* (1992) demonstrated that infection of mango fruits before harvest occurred through endophytic colonization of the pedicel tissues by *Botryosphaeria* spp. present from previous growth flushes. The possibility of pruning to promote new growth flush was tested as a means to reduce inoculum in the stem tissue from which new-season inflorescences emerged. Cooke *et al.* (1998) reported that the levels of endophytic organisms such as *Botryosphaeria* spp. were reduced significantly when a pruning programme was implemented in mango orchards as a preharvest control measure. Korsten (2006) found that prevention of water stress during fruit development and maturation, and avoidance of placing fruits on the ground suppressed disease development. He also suggested that fruits should be cooled to 13°C immediately after harvest and stored in a well-ventilated place.

Physical control

Growing public demand for chemical residue-free fruits has encouraged the development of alternative technologies, such as irradiation, heat treatment and cold-temperature storage. Cold storage of mango fruit (10–12°C) is one of the best ways of delaying fruit ripening and, thus, decreasing post-harvest decay (Sivankalyani *et al.*, 2016). Shortwave infrared radiation treatments reduce anthracnose damage in mango (Saaiman, 1996) and this approach can also be considered for the organic market.

Heat treatment is known to reduce post-harvest diseases. Different approaches have been used, such as hot-water dipping and rinsing, and hot water vapour and dry-air treatments (Schirra *et al.*, 2000). There are many benefits to heat treatments, such as reduction in post-harvest decay, killing of pests, colour and flavour preservation and shelf-life improvement, among others (Lurie, 1998; Schirra, 2000; Fallik, 2004). Hot-water brushing at 50–60°C for 20 seconds after harvesting reduces decay development via both surface cleansing and induction of fruit resistance against pathogens (Prusky *et al.*, 1996a; Fallik, 2004); this method is applied in Israel. Hot-water dipping for 3–7 minutes has been recommended and is moderately efficient at delaying post-harvest rot (Johnson, 1994). Hot-water dips, or spray can control fungal infections such as anthracnose and alternaria rot better than stem-end-rot (Johnson, 1994). Trials using gamma irradiation to control mango anthracnose have concluded that incorporation of hot fungicide dip is necessary to improve disease control afforded by irradiation (Chadha, 1989).

Regarding stem-end rot, for high-value fruit, especially those destined for export, various post-harvest treatments have been beneficial (Ploetz, 2018). For example, Alvindia and Acda (2015) reported a 48–61% reduction in stem-end rot of 'Carabao' fruits after 20 min in 53°C water. Terao *et al.* (2015) indicated that a low dose (< 3 kJ m⁻²) of UV-C irradiation helped manage post-harvest diseases of mango caused by *B. dothidea*, *L. theobromae*, *A. alternata* and *C. gloeosporioides*, even though a direct impact on the pathogens was not evident. Santos *et al.* (2015) suggested that a dose of 0.45 kGy of gamma irradiation reduced disease caused by *L. theobromae*.

Fruit sanitizers

The purpose of fruit sanitizers is to wash and kill the microorganisms on the fruit surface. Traditionally, the sanitizers consisted of water with or without chemicals. One of the most extensively used and studied sanitizers is chlorine (water pH 6.5–7.5; chlorine concentration 100–150 ppm). In addition to chlorine, sulphur dioxide has also been used as fungal disinfectant (Johnson *et al.*, 1997; Tefera *et al.*, 2007). Different forms of chlorine, such as sodium hypochlorite, calcium hypochlorite and chlorine gas, control a wide range of post-harvest pathogens (Boyette, 1995). In the past, elevated chlorine dosages were frequently used due to the misconception that chlorine leaves no residues on the fruit. Common alternatives to chlorine are ozone (O₃), oxidized water and hydrogen peroxide. Ozone and ozonated water were recognized in 1997 by the FDA as safe food disinfectants and were proven to control post-harvest rots of mango (Monaco *et al.*, 2014). Recently, ozonated water has been reported as a sanitizer for mango cv. Palmer as it increases antioxidant activity (Minas *et al.*, 2012; Lima *et al.*, 2014, Monaco *et al.*, 2016).

Electrolyzed water has also been suggested as a sanitizer for the industry (Colangelo *et al.*, 2015). Electrolyzed water is produced by adding sodium chloride (as an electrolyte) to tap water and passing an electrical current through an anode or cathode to produce oxidizing (acidic) and reducing water (alkaline), respectively. The high electrolyzed water potential works against both bacteria (Pinto *et al.*, 2015) and fungi (Guentzel *et al.*, 2010). Hydrogen peroxide has also been recommended as an effective disinfectant against several fungi (Boyette, 1995).

BACTERIAL BLACK SPOT (BACTERIAL CANKER)

Resistance to Bacterial Black Spot varies greatly among mango cultivars, and resistant cultivars should be used where disease pressure is high (Manicom and Pruvost, 1994). Pathogen-free planting material should be utilized when new orchards are established. The pathogen moves only short distances in wind-blown aerosols (usually within orchards) (Gagnevin and Pruvost, 2001), and long-distance dissemination occurs almost entirely via infected propagation material and less frequently in surface-contaminated seeds (Manicom and Pruvost, 1994). Windbreaks should be used to reduce wounding and infected twigs should be removed from the canopy.

FOLIAR AND FLORAL DISEASES

SECA AND SUDDEN DECLINE

Given the destructive impact of these diseases, preventing the dissemination of these pathogens to new areas should be a high priority (Ploetz, 2018). Pathogen-free propagation material is needed whenever new plantings are established and germplasm is moved. Clean pruning tools should be used in affected areas and should be frequently disinfested with bleach, formalin or other disinfectants (Junqueira *et al.*, 2002). Trees that have been killed by the disease should be removed and destroyed because they are significant reservoirs of the vector and pathogens.

The use of resistant or tolerate genotypes is an appropriate alternative. Various levels of tolerance have been observed and resistant materials have been developed. However, pathogenic variation in the causal fungus in Brazil has hindered progress (Rossetto *et al.*, 1996; Junqueira *et al.*, 2002; Silveira *et al.*, 2006). In Brazil, the disease responses of some genotypes vary in different production areas, but 'Manga Dagua', 'Pico', 'IAC 101', 'IAC 102', 'Edwards', 'Van Dyke' and 'Carabao' show resistance to two pathotypes, while 'Rosa', 'Sabina', 'Sao Quirino', 'Oliveira Neto', 'Jasmim', 'Sensation', 'Irwin' and 'Tommy Atkins' are generally tolerant (Ribiero, 1997; Junqueira *et al.*, 2002). 'Espada' is also reported to be tolerant, but old trees are attacked. Colosimo *et al.* (2000) reported that 'Oliveira' was the most resistant, while 'Carlota', 'Imperial', 'Extrema' and 'Pahiri' had intermediate resistance. Carvalho *et al.* (2004) described two new cultivars, 'IAC 103 Espada Vermelha' and 'IAC 109 Votupa', with moderate resistance to seca. 'IAC 103 Espada Vermelha' also had moderate resistance to powdery mildew but was susceptible to anthracnose, and both cultivars were susceptible to malformation.

In Oman, 'Hindi Besennara', 'Sherokerzam', 'Mulgoa', 'Baneshan', 'Rose' and 'Alumpur Baneshan' developed significantly less disease when challenged with *Ceratocystis maginecans* (Al Adawi, *et al.*, 2013).

MALFORMATION

Management of malformation can be difficult and mainly relies on proper sanitation and hygiene in the orchards. Thus, new plantings should be established with pathogen-free nursery stock and, consequently, scion material should never be taken from affected orchards, and any affected plants observed in the nursery should be removed and burned. Once the disease is found in an orchard, all affected terminals and the subtending three nodes must be removed from the trees and burned (Prusky *et al.*, 2009). Good results have been observed following this recommendation, as those reported by Schoeman *et al.* (2018), who found that continuous removal of infected inflorescences over a four-year period resulted in a significant decline of disease incidence. However, it may be difficult to apply this treatment on large trees (Prusky *et al.*, 2009).

The selection of resistant varieties can play a vital role in reducing the malformation incidence, especially in the most severely affected areas (Katoch *et al.*, 2019). Variation in the susceptibility

to malformation of different varieties has been reported (Prakash and Srivistava, 1987; Fayyaz *et al.*, 2002). In some cases, results have been inconsistent (Ploetz, 2001), but some local cultivars that have consistently been deemed resistant, such as 'Bhadauran' in Brazil and India, 'Primor' in Brazil and 'Zebda' in Egypt, should receive wider attention (Ploetz, 2018).

POWDERY MILDEW

Reduction of inoculum potential of the pathogen at early stages is likely to decrease disease incidence (Joubert, 1991). Regular inspection of mango orchards and removal/pruning of infected leaves and malformed panicles reduce the load of primary inoculum and improve fungicidal control (Prakash and Misra, 1992, 1993a; 1993b; Prakash and Raof, 1994).

Mango cultivars vary in their resistance to powdery mildew (Palti *et al.*, 1974). 'Zill', 'Kent', 'Alphonso', 'Seddek' and 'Nam Doc Mai' are very susceptible; 'Haden', 'Glenn', 'Carrie', 'Zebda', 'Hindi be Sennara', 'Ewaise' and 'Keitt' are moderately susceptible; and 'Sensation', 'Tommy Atkins' and 'Kensington' are slightly susceptible (Ploetz *et al.*, 1994; Nofal and Haggag, 2006). In India, Tiwari *et al.* (2006) reported that 'Baigan Phalli', 'Barbalia', 'Dabari', 'Dilpasand', 'Khirama', 'Nagarideeh', 'Oloor' and 'Totapari' were highly resistant and 'Amrapali' was most susceptible.

3. CAN AGROHOMEOPATHY BE USED TO EFFICIENTLY MANAGE OR CONTROL PESTS AND DISEASES IN MANGO?

Homeopathy is based on the principle that a substance which in a massive dose generates pathological symptomatology has the possibility to cure a disease, if applied in the minimum doses obtained by dilution and intense agitation. This notion was proposed in 1796 by Samuel Hahnemann (Toledo *et al.*, 2011). Thus, the central tenet of homeopathy is that "like cures like". Homeopathic remedies are prepared ('potentized' or 'dynamized') in steps of alternately diluting and succussing a homeopathic stock, known as 'mother tincture', until they reach calculatory dilutions beyond Avogadro's number, implying a non-molecular action of remedies with specific healing properties. One of the most paradigmatic and controversial aspects of homeopathy is if these serial dilutions would have a measurable effect even when they are given to an animal or a plant in infinitesimal doses (Mazón-Suastegui *et al.*, 2019), although the Swiss chemist Louis Rey found that the structure of hydrogen bonds in homeopathic dilutions of salt solutions is very different from that in pure water, probably due to vigorous shaking of solutions that takes place during homeopathic 'succussion' (Rey, 2003). Products labeled as homeopathic and currently marketed in the U.S. have not been reviewed by the FDA (USA Food & Drugs Administration) for safety and effectiveness to diagnose, treat, cure, prevent or mitigate any diseases or conditions, and, therefore, there are no FDA-approved products labeled as homeopathic (FDA, 2020). In fact, homeopathy is considered by the scientific community as a pseudoscientific system of alternative medicine (Smith, 2012).

Agrohomeopathy, or the application of homeopathic principles and remedies to agriculture, has drawn emerging interest over the last years, since it could represent an ecological alternative to decrease or eliminate agrochemical use (Abasolo-Pacheco *et al.* 2020). However, no reports on this practice can be found in the main horticultural journals and, consequently, the results reported so far in more obscure journals should be taken with caution.

Homeopathic preparations have been described as relatively cheap, with few or no ecological side-effects, and seemingly harmless, which would make this discipline optimally suited to the

holistic approaches of organic and, above all, biodynamic agriculture (Carpenter-Boggs *et al.*, 2000). According to Bonato (2007), the use of high dilution preparations following homeopathy could work on plants by improving their physiological status, inducing resistance and favouring multiple biological interactions in the way that healthy plants are tolerant to pests and diseases and promptly response for optimal production (Bonato, 2007). Positive results from the use of homeopathic medicines on different plant species have been claimed in seed germination, emergence and initial growth (Abasolo-Pacheco *et al.*, 2020), in overall production (Rossi *et al.*, 2007), in remediation of soils affected by heavy metals (Dos Santos *et al.*, 2016), in mitigation of the effects of abiotic stress, such as salinity (Giardini-Bonfim *et al.*, 2012) or hot and dry weather (Pelikan and Hunger, 1971), and to control plant pests and diseases (Modolon *et al.*, 2012; Ramaia & Kumar, 2015). Studies on plant diseases control using homeopathic preparations report experiments based on phytopathological models (plants naturally infected or artificially inoculated with fungi, viruses, bacteria, nematodes), in vitro spore germination and growth models, and field trials (agronomical and phytopathological experimentations) (Betti *et al.*, 2009). In fruit crops, experimental effort has mainly focused on fruit rot diseases control. For instance, Baviskar and Suryawanshi (2015) found that *Arsenicum album* had an inhibitory activity against fungi responsible for fruit rot diseases in apple. Kehri and Chandra (1986) evaluated some homeopathic treatments against *Lasiodiplodia theobromae*, (Pat.) Griffon & Maubl., a severe pathogen that causes postharvest rot of guava (*Psidium guajava* L.), and reported that all the tested potencies of *Arsenicum album* completely suppressed in vitro spore germination and, in addition, largely reduced fruit rots when applied as a pre-inoculation dip treatment of guava fruits. Yadav *et al.* (2013) reported good results with Methyl jasmonate 200 for Fusarium fruit rot of banana.

Specific studies on the use of agrohomoepathic approaches in mango are scarce. Khanna and Chandra (1978) determined the fungitoxicity of 10 homeopathic drugs against *Pestalotia mangiferae* Henn., the causal agent of mango fruit rot in India, in terms of the inhibition of spore germination. To do this, spores of the pathogen were suspended in different potencies (1-200) of the drugs and the hanging drop technique of Hoffman (Hoffman, 1869) was employed. The drugs that completely inhibited spore germination after 8-12 hour-time incubation were screened for their efficacy in controlling fruit rot. For this purpose, healthy freshly picked mango fruits of cv. 'Dusehri' were injured with a sterilized needle, artificially inoculated with a spore suspension, and dipped, in pre- and post-inoculation treatments, for 3-5 min. in each drug or sterilized distilled water (control). The authors claimed that, although several drugs were able to completely inhibit spore germination of the pathogen, only *Lycopodium clavatum* potency 190 was effective in reducing fruit infection percentage and rot percentage after 8 days, and thus, they suggested that it may be safely recommended for the control of mango fruit rot caused by *P. mangiferae*.

More recently, Alam *et al.* (2017) tested 72 homeopathic solutions containing anti-fungal properties, each with 5 Centesimal Hahnemannian potencies, to control mango anthracnose caused by *Colletotrichum gloeosporioides* (Penz. & Sacc.). They made two types of experiments. One, using Petri dishes, where the solutions were tested individually against mycelial growth of *C. gloeosporioides* using Oat Meal Agar (OMA) medium by food poisoning method. Another, with recently harvested fruits from three cultivars treated by dipping for 10 minutes in a solution with the best solutions selected in the Petri dish experiment. The authors claimed that the results from the in vitro study pointed out that all the homeopathic drugs showed remarkable efficacy for inhibition of *Colletotrichum gloeosporioides* mycelia growth, especially *Arsenicum album* Q at 10000 ppm, with a 96.40% inhibition percent. *Arsenicum album* Q at 10000 ppm concentration was used in the fruits experiment, and eighteen days after treating inhibition of Percent Disease Incidence was different in each cultivar: 18.2% in cv. 'Himsagar', 8% in cv. 'Amrapali' and 0% in cv. 'Langra', although lesion size and lesion cover were significantly

reduced by the homeopathic treatment in all three cultivars. The authors concluded that these results may suggest the potential of homeopathic medicines for controlling anthracnose of mango and other fruits.

Although these results could seem to be promising and agrohomoepathy has been proposed as an alternative to the widespread use of herbicides, pesticides and other chemicals in agriculture, basic and applied research on effectiveness of homeopathic treatments in crops is scarce and does not often provide sufficient information to be interpreted properly. No significant works in this field are found in relevant horticultural journals and, consequently, caution should be taken on the possible effects of these treatments in agriculture. A more appropriate approach to reduce the use of chemical treatments could probably be the use of organic agriculture.

4. WHAT KIND OF ALTERNATIVE CAN THE KNF (KOREAN NATURAL FARMING) PRESENT AS POTENTIAL STRATEGY TO CONTROL OR MANAGE PESTS AND DISEASES IN MANGO?

Basic principles of Korean Natural Farming (KNF) include using indigenous microorganisms (IMO), maximize the potential of natural environment, minimize the use of synthetic fertilizers, practice no till, eliminate emission of livestock waste effluents and increase production with less inputs (Wang *et al.*, 2012). Therefore, this farming approach, developed by Han Kyu Cho from Janong Natural Farming Institute in South Korea (Cho and Cho, 2010), maximizes the use of on-farm resources, recycles farm waste, and minimizes external inputs while fostering soil health (Wang *et al.*, 2012). KNF has been practiced for decades in Asia and was introduced to Hawaii at the end of 1990s, where it has been gaining popularity among farmers who are interested in sustainable agriculture (Wang *et al.* 2012).

Cultivation of IMO from the farm instead of introducing alien beneficial organisms is a key practice of KNF (Wang *et al.*, 2012). This involves a four-step process of capturing, cultivating, and preserving IMO, to create products that are often referred to as IMO 1, IMO 2, IMO 3, and IMO 4 (Keliikuli *et al.*, 2019). A list of natural inputs used to make the IMOs and a thorough description of how to prepare them can be found in Park and Duponte (2008) and Keliikuli *et al.* (2019). The final product, IMO 4, should be used as a top dressing, gently mixing 150 kg/0.1 ha into the topsoil, and covering the inoculated soil with mulch (i.e., bamboo leaf litter, wood chips, etc.) (Keiikuli *et al.*, 2019). It is recommended that IMO 4 be applied to the soil seven days before seeding or transplanting and two to three hours prior to sunset, since treating the soil in the late afternoon gives the microbes more time to adjust to the environmental changes, particularly the increase in temperature (Cho and Cho 2010).

Foliar sprays of composts made from various herbs or farm wastes is another key practice of KNF (Wang *et al.*, 2012). According to these authors, the application of nutrient sprays may be environmentally friendly since the nutrients are directly delivered to the plant in limited amounts, and involves benefits to young seedlings with smaller root systems, reductions in the amount of N application, better nutrient uptake during the reproductive stage due to a decrease in root activity, and the ability to modify the nutrient inputs accordingly.

Studies in Hawaii, where farmers' plots for vegetable production using KNF practices were compared to control plots using conventional or organic farming, found that KNF did significantly improved some conditions of plant and soil health, especially in plots with soybean, and required

less irrigation than conventional farming, probably due to the soil mulching required by KNF (Wang *et al.*, 2012). Nevertheless, KNF did not protect plants from insects and diseases.

Scientific evidence of the benefits of KNF has been limited (Wang *et al.*, 2012), and, as far as we know, there are no references about the effects of KNF on control of fruit crops diseases. Probably, the approach closest to IMO, one of the foundations of KNF, could be the use of indigenous biocontrol agents (BCAs) as a tool to reduce or exterminate pathogen populations (AbuQamar *et al.*, 2017; Syed Ab Rahman *et al.*, 2018). This is since epiphytic microbes that live on plant surfaces without causing any symptoms to the plants are known for their efficacy to inhibit plant pathogens (McGrath and Andrews 2005; Janisiewicz *et al.* 2010; Janisiewicz *et al.* 2013). We have found some references of disease control with BCAs in mango. For instance, In Mexico, Bautista-Rosales *et al.* (2014) evaluated the effectiveness of the antifungal yeast *Cryptococcus laurentii* [(Kuff.) C.E. Skinner] strain LSD, isolated from the surface of local 'Ataulfo' mango fruits, against the causal agent of anthracnose, *Colletotrichum gloeosporioides* ((Penz.) Penz. & Sacc.) and they found that *C. laurentii* showed a high antagonistic potential *in vivo*, with significant inhibition of anthracnose (75.88%), probably due to competition for nutrients and for space. In Thailand, Rungjindama (2016) reported that two isolates of epiphytic microbes (bacteria and yeasts) from healthy leaves and fruits of local mangoes reduced lesion sizes caused by *C. gloeosporioides* compared to control treatment. In the United Arab Emirates (UAE), Kamil *et al.* (2018) found that 19 actinobacterial isolates obtained from mango rhizosphere soils in the UAE showed antagonistic activities against *Lasiodiplodia theobromae* (Pat.) Griffon and Maubl., causal agent of mango dieback disease in different areas of the world, including Brazil, Korea, India, Oman, Pakistan, USA and the UAE. All those isolates were screened *in vivo* for their abilities to reduce lesion severity on fruits inoculated with *L. theobromae*, and the three isolates showing the strongest inhibitory effect against this pathogen were inoculated on mango seedlings, 1 week before inoculation with *L. theobromae*, and exhibited very high levels of disease protection against the pathogen. According to these authors, the indigenous strains identified in this study were safe, inexpensive, long lasting, and well-suited to extreme harsh conditions, which make them suitable to be incorporated into sustainable IPM strategies to manage dieback disease in mango orchards in different countries.

According to these promising results, it seems that the use of indigenous BCAs, alone or combined with some practices including in KNF to increase plant and soil health (such as soil mulching, foliar sprays with core solutions, etc.), could be potential strategies to manage diseases of mango. In fact, some principles of KNF, such as enhancing soil microbial activity and fertility, increasing microbial diversity or maximizing the use of on-farm resources, are also found in other types of organic farming and, consequently, organic mango production could be an interesting way for pest and disease control. This is an area where additional work is clearly needed and should be included in a holistic organic management.

5. FUTURE AREAS AND LINES OF RESEARCH FOR THE CONTROL OF MANGO PESTS AND DISEASES

Significant advances have been made in the last decades in optimizing the control of different pests and diseases combining chemical, biological and management approaches. However, new pests and diseases are constantly appearing in different countries and globalization is helping the spread of pests and diseases to new areas. Consequently, internationally coordinated efforts are required to optimize pest and disease control along different areas and lines of research. A holistic approach combining biological and chemical control, cultural practices, and plant selection and breeding should be combined. Examples of lines of research where information are still needed include the following:

- Biological control: research is needed on identifying additional pest and disease native natural enemies and on optimizing their populations through appropriate orchard management to minimize damage.
- Cultural practices: agroecology is a field in which additional research is needed including different topics such as ecological landscaping, crop rotations, use of cover crops, optimize water use or developing strategies to improve pest and disease monitoring. In this sense, new technologies (drones, 5G) can be useful for early damage identification.
- Chemical control: minimize the use of synthetic products and increase the range of available naturally occurring biopesticides (pheromones, insect growth regulators, etc.).
- Genetics and diversity: selection and development of tolerant and resistant varieties including the use of biotechnological approaches (genetics, genomics, in vitro culture).
- Appropriate control of imports of fruits and plant material, including periods under quarantine if necessary, should be prioritized with fast protocols for pest and diseases detection.
- Increase the investment for research in developing countries where most of the mango production is based and resources devoted for pest and disease control are scarce.
- Studies are needed on the effect of climate change on pest and disease control and management.

6. References

- Abasolo-Pacheco, F., Bonilla-Montalván, B., Bermeo-Toledo, C., Ferrer-Sánchez, Y., Ramírez-Castillo, A.J., Mesa-Zavala, E., Llerena-Ramos, L. & Mazón-Suástegui, J.M., 2020. Efecto de medicamentos homeopáticos en plantas de tomate (*Solanum lycopersicum* L.). *Terra Latinoamericana* Número Especial 38-1: 219-233.
- Abdel Kareim, A.L., Ragab, M.E., Ghanim, M.N. & Abd El-Salam, S.A., 2018. Seasonal activity, natural enemies and life table parameters of *Cryptoblabes gnidiella* Mill. on mango inflorescences. *J. Plant Prot. and Path., Mansoura Univ.*, Vol.9 (7): 393 – 397.
- Abdulla, N., Rwegasiri, G., Jensen, K., Mwatalala, M. & Offenber, J., 2016. Control of mango seed weevil (*Sternonchetus mangiferae*) using the African weaver ant (*Oecophylla longinoda* Latreille) (Hymenoptera: Formicidae). *Journal of Applied Entomology*, 140: 500–506.
- AbuQamar, S. F., Moustafa, K., & Tran, L. S., 2017. Mechanisms and strategies of plant defense against *Botrytis cinerea*. *Crit. Rev. Biotechnol.* 3: 262–274. doi: 10.1080/07388551.2016.1271767.
- Acuna, H.E. & Waite, B.H., 1977. La muerte regresiva del mango (*Mangifera indica* L.) en El Salvador. *Proceedings of the American Society of Horticultural Sciences, Tropical Region* 21: 15–16.
- Ahmed, W., Nawaz, M.A., Saleem, B.A. & Asim, M., 2005. Incidence of mango midge and its control in different mango growing countries of the world. First International Conference on Mango and Date palm, At Institute of Horticultural Sciences University of Agriculture, Faisalabad- Pakistan.
- Akem, C. N., 2006. Mango anthracnose disease: Present status and future research priorities. *Plant Pathology Journal* 5: 266–73.
- Al Adawi, A. O., Deadman, M. L., Al Rawahi, A. K., Al Maqbali, Y. M., Al Jahwari, A. A., Al Saadi, B. A., Al Amri, I. S. & M. J. Wingfield., 2006. Aetiology and causal agents of mango sudden decline disease in the Sultanate of Oman. *European Journal of Plant Pathology* 116: 247–54.
- Al Adawi, A. O., Al Sadi, B. A., Deadman, M. L., Barnes, I., Al Jabri, M. H., Wingfield, B. D. & Wingfield, M. J., 2013. Evaluation of mango cultivars for resistance to infection by *Ceratocystis manginecans*. *Acta Horticulturae* 992: 393–406.
- Al Subhi, A. M., Al Adawi, A. O., Van Wyk, M., Deadman, M. L. & Wingfield, M. J., 2006. *Ceratocystis omanensis*, a new species from diseased mango trees in Oman. *Mycological Research* 110: 237–245.
- Alam, A., Adhikary, S.K. & Ahmed, M., 2017. Effectiveness of selected homeopathic medicines against *Colletotrichum gloeosporioides*. *Asian J. Plant Pathol.*, 11: 118-129.
- Almeida, R.T. de, Landim, C.M.U. & Teixeira, L.M.S., 1979. Seedling blight (*Sclerotium rolfsii* Sacc.) of cashew and mango in the state of Ceara, Brazil. *Fitossanidade* 3: 40–41.
- Aluja, M., 1993. “Manejo Integrado de la Mosca de la Fruta”. Editorial Trillas, Mexico Distrito Federal, Mexico.
- Aluja, M., 1996. Future trends in fruit fly management. In: “Fruit Fly Pests: a World Assessment of their Biology and Management”, edited by McPherson, B. and Steck, G. Published by St Lucie Press, Delray Beach, Florida. Pp. 309–320.
- Aluja, M. & Mangan, R.L., 2008. Host status determination for pestiferous fruit fly (Diptera: Tephritidae) species: conceptual, methodological and regulatory considerations. *Annual Review of Entomology* 53: 473–502.

Aluja, M. & Pinero, J., 2004. Testing human urine as a low-tech bait for *Anastrepha* spp. (Diptera: Tephritidae) in small guava, mango, sapodilla and grapefruit orchards. *Florida Entomologist* 87: 41–50.

Aluja, M., Cabrera, M., Guillen, J., Celedonio, H. & Ayora, F., 1989. Behavior of *Anastrepha ludens*, *A. obliqua* and *A. serpentina* (Diptera: Tephritidae) on a wild mango tree (*Mangifera indica*) harbouring three McPhail traps. *Insect Science and its Application* 10: 309–318.

Aluja, M., Sivinski, J., Ovruski, S., Guillen, L., Lopez, M., Cancino, J., Torres-Anaya, A., Gallegos-Chan, G. & Ruiz, L., 2009. Colonization and domestication of seven species of native New World Hymenopterous larval-prepupal and pupal fruit fly (Diptera: Tephritidae) parasitoids. *Biocontrol Science and Technology* 19 (Suppl. nº 1): 49-79.

Alvarez-Garcia, L.A. & Lopez-Gracia, J., 1971. Gummosis, dieback, and fruit rot disease of mango (*Mangifera indica* L.) caused by *Physalospora rhodina* (B. and C.) Cke. in Puerto Rico. *Journal of Agriculture University of Puerto Rico* 55: 435–450.

Alvindia, D. G. & Acda, M. A., 2015. Revisiting the efficacy of hot water treatment in managing anthracnose and stem-end rot diseases of mango cv. 'Carabao'. *Crop Protection* 67: 96–101.

Angulo, S. M. & Villapudua, J. R., 1982. Buba of mango (*Mangifera indica* L.) in the state of Sinaloa, Mexico. *Phytopathology* 72: 171.

Anonymous, 1984. Major insect pests and diseases of mango and their control. In: "The Mango Technical committee. The Philippines recommends for Mango". Philippines Agriculture and Resources Research Foundation. 124 pp.

Anonymous, 1987. Red-banded mango caterpillar *Noorda albizonalis Hampsoni* (Lepidoptera: Pyralidae). *Plant Quarantine Leaflet-Australia* 51.

Anonymous, 1989. Mango Pests and Disorders. Information Series QI89007. Queensland Department of Primary Industries, Brisbane, Australia.

Anonymous, 2002. Fruit flies eradicated from Nauru. 3pp. Available at: http://www.pacifly.org/Success_stories/Nauru_eradl.htm (accessed 10 September 2008).

Anonymous, 2005. Malathion (CAS Number 121-75-5). Toxicology information about insecticides used for eradicating mosquitoes. Available at: <http://www.atsdr.cdc.gov/consultations/malathion.html> (accessed 20 July 2007).

Anonymous, 2006. Importation of Fresh Mango Fruit (*Mangifera indica*) from India into the United States, a Qualitative Pest Risk Assessment. Animal and Plant Health Inspection Service, Plant Protection Quarantine, Raleigh, North Carolina.

Arias, M. & Jines, A., 2004. *Manejo Integrado de Moscas de la Fruta en el Litoral Ecuatoriano*. Instituto Nacional de Investigaciones Agropecuarias – Programa de Modernización de Servicios Agropecuarios, Guayaquil, Ecuador.

Arias, M., Jines, A., Bustos, P., Pluas, M. & Carrera, C., 2004a. Enemigos Naturales de *Aulacaspis tubercularis* (Homoptera: Diaspididae) en Mango. Instituto Nacional Autonomo de Investigaciones Agropecuarias (INIAP), Boliche, Guayas, Ecuador.

Arias, M., Jines, A., Maldonado, E. and Pluas, M., 2004b. *Cybocephalus nipponicus* Endrody-Younga (Coleoptera: Nitidulidae) un predator exótico para el control de *Aulacaspis tubercularis* en mango de exportación. Instituto Nacional Autonomo de Investigaciones Agropecuarias (INIAP), Boliche, Guayas, Ecuador

Assis, E. & Rabelo, F., 2005. Pragas da Mangueira, Monitoramento, Nivel de Acao e Controle. Empresa Brasileira de Pesquisa Agropecuaria, Embrapa Semi-Arido, Ministerio da Agricultura, Pecuaria e Abastecimento, Petrolina, Brazil.

Arauz, L.F., 1999. Optimizacion economica del combate de enfermedades en cultivos: un ejemplo basado en la antracnosis del fruto del mango. (Abstract). In: *XI Congreso Nacional Agron. Recursos Naturales, V Congreso Nacional Entomol., IV Congreso Nacional Fitopatol., III Congreso Nacional Suelos I Congreso Nacional Ext. Agric. Forestal*. Vol. II. San Jose, Costa Rica. Pp. 61.

Arauz, L.F., 2000. Mango anthracnose: economic impact and current options for integrated and management. *Plant Disease* 84 (6): 600–611.

Arjona-Girona, I & López-Herrera, C.J., 2018. First report of *Rosellinia necatrix* causing White root rot in mango trees in Spain. *Plant Disease* 102 (12): 2639.

Atkinson, T.H., Foltz, J.L., Wilkinson, R.C. & Mizell, R.F., 2017. Granulate ambrosia beetle *Xylosandrus crassiusculus* (Motschulsky) (Coleoptera: Curculionidae: Scolytinae). EENY 131. Department of Entomology and Nematology, UF/ Institute of Food and Agricultural Sciences (IFAS) Extension, University of Florida.

Azmy, A.M.K., 2014. Controlling of Mango Powdery Mildew by some Salts, Growth Regulators and the Biofungicide AQ10 Compared with Punch Fungicide in Egypt, *American Journal of Life Sciences*. Special Issue: *Role of Combination Between Bioagents and Solarization on Management of Crown-and Stem-Rot of Egyptian Clover*. Vol. 2, No. 6-2: 33-38.

Bajracharya, A.S.R., Piya, S. & Ghimire, K., 2012. Effectiveness of chemical insecticides against mango stone weevil (*Sternochetus mangiferae*; Fabricius; Coleopter: Curculionidae) in the foothills of Eastern Nepal. Proceedings of the 10th National Outreach Research Workshop, At Regional Agricultural Research Station, Lumle, Kaski, Nepal. Pp.: 256-259.

Balock, J.W. & Kozuma, T., 1964. Notes on the biology and economic importance of the mango weevil, *Sternochetus mangiferae* (Fabricius), in Hawaii (Coleoptera: Curculionidae). *Proceedings of the Hawaiian Entomological Society* 8: 353–364.

Balock, J.W. & Lopez, D.F., 1969. Trapping for control of the Mexican fruit fly in mango and citrus groves. *Journal of Economic Entomology* 62: 53–56.

Barbosa, F.R., 2005. Arthropodes-pragas associados a cultura da mangueira no Brasil e seu controle. In: “Pragas da Mangueira, Embrapa Semi-arido”, edited by Menezes, E. and Barbosa, F.R. Ministerio da Agricultura, Pecuaria e Abastecimento, Petrolina, Brazil. Pp. 1–50.

Batista, A.C., 1947. Mal do recife (grave doenca da mangueira). Unpublished thesis, Pernambuco College of Agriculture, Pernambuco, Brazil (Abstract in *Review of Applied Mycology*, 1948, 27: 77–78).

Batzer, J.C., Gleason, M.L., Harrington, T.C. & Tiffany, L.H., 2005. Expansion of the sooty blotch and fly speck complex on apples based on analysis of ribosomal DNA gene sequences and morphology. *Mycologia* 9: 1268–1286. *Proceedings of the Hawaiian Entomology Society* 27: 367–378.

Bautista-Rosales, P.U., Calderon-Santoyo, M., Servín-Villegas, R., Ochoa-Alvarez, N.A., Velazquez-Juarez, R. & Ragazzo-Sanchez, J.A., 2014. Biocontrol action mechanisms of *Cryptococcus laurentii* on *Colletotrichum gloeosporioides* of mango. *Crop protection* 65: 194-201.

Baviskar, R.N. & N.S. Suryawanshi, 2015. Application of certain homoeopathic medicines used against fruit rot of apple caused by *Penicillium expansum* Link. *Int. J. Life Sci.* 3: 96-98.

Berthet, J. A., 1914. Molestia da mangueira. *Bolm Agric. (Sao Paolo)* 15: 818–19.

- Bess, H.A., van den Bosch, R. & Haramoto, F.H., 1961. Fruit fly parasites and their activities in Hawaii.
- Betti, L., Grazia, G., Majewsky, V., Scherr, C., Shah-Rossi, D., Jäger, T. & Baumgartner, S., 2009. Use of homeopathic preparations in phytopathological models and in field trials: a critical review. *Homeopathy* 98: 244–266.
- Bitancourt, A.A. & Jenkins, A.E., 1943. Scab of mango caused by *Elsinoe*. (Abstract). *Phytopathology* 33: 1.
- Bonato, C.M., 2007. Homeopatia em modelos vegetais. *Cultura Homeopática* 21: 24-28.
- Bouagga, S., Hassan, N., Ben Halima, M., Jammazi, A., Djelouah, K. & Al-Zaidi, S., 2004. Evaluation of an “Attract and Kill” System to Combat *Ceratitis capitata* on Peach Trees in Tunisia. *Journal of Agricultural Science and Technology* 4: 612-619.
- Boyette, M. D. (1995). Packaging Requirements for Fruits and Vegetables. In: “Postharvest Handling and Cooling of Fresh fruits, Vegetables and Flowers for Small Farm 800–4”. North Carolina Crop Extension Services. USA. Pp. 17.
- Brent, K. J. & Hollomon, D. W., 2007. *Fungicide Resistance: The Assessment of Risk*. Fungicide Resistance Action Committee. Croplife International: Brussels, Belgium.
- Britz, H., Steenkamp, E. T., Coutinho, T. A., Wingfield, B. D., Marasas, W. F. O. & Wingfield, M. J., 2002. Two new species of *Fusarium* section *Liseola* associated with mango malformation. *Mycologia* 94: 722–30.
- Burns, R.E., Harris, D.L., Moreno, D.S. & Eger, J.E., 2001. Efficacy of spinosad bait sprays to control Mediterranean and Caribbean fruit flies (Diptera: Tephritidae) in commercial citrus in Florida. *Florida Entomologist* 84: 672–678.
- Burrack, H.J., Connell, J.H. & Zalom, F.G., 2008. Comparison of olive fruit fly (*Bactrocera oleae* (Gmelin)) (Diptera: Tephritidae) captures in several commercial traps in California. *International Journal of Pest Management* 54: 227–234.
- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J., 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology* 97: 286–292.
- Butani, D.K., 1979. Insects and fruits. Periodical Expert Book Agency, Delhi.
- CABI, 2009. Mango midges. Survey of midges and their natural enemies associated with mango and develop non pesticides measures for their control in Pakistan. <https://www.cabi.org/uploads/projectsdb/documents/6049/Mango%20Midges%20in%20Pakistan,%20June%202009.pdf>.
- CABI, 2020. *Xylosandrus crassiusculus* (Asian ambrosia beetle). Datasheet. Retrieved from <https://www.cabi.org/isc/datasheet/57235#6CE9E9B6-8C1E-4666-A0C2-B8BFF8FB7EF2>.
- Cabrera, H., Villanueva, J. and Ortega, D. (1993) Fruit fly infestation timing on ‘Manila’ mango fruit at Veracruz, Mexico. (Abstract). *IV International Mango Symposium*, p. 118.
- Cabrera-Mireles, H. & Ortega-Zaleta, D. A., 2004. Principales plagas del mango. In: “Tecnología para la producción forzada de mango ‘Manila’ con calidad fitosanitaria en Veracruz”, edited by INIFAP-CP-Veracruz. INIFAP-CIRGOC-Campo Experimental Cotaxtla. Memoria Técnica 15. División Agrícola, Veracruz, Mexico. Pp.: 13-31.
- Campbell, R.J. (ed.), 1992. *A Guide to Mangoes in Florida*. Fairchild Tropical Garden, Miami, Florida.

- Carpenter-Boggs, L., Reganold, J.P. & Kennedy, A.C., 2000. Biodynamic preparations; short-term effects on crops, soils and weed populations. *Am. J. Alternative Agr.* 15: 110–118.
- Carvalho, P.J. & De Queiroz, A.C. (eds), 2002. *A Cultura da Mangueira*. CIP-Brasil, Brasilia.
- Carvalho, R. da S., Nascimento, A., Morgante, J. & Fonseca, N., 1996. Susceptibility of different mango varieties to the attack of the fruit fly, *Anastrepha obliqua*. In: "Fruit Fly Pests: a World Assessment of their Biology and Management", edited by McPherson, B. and Steck, G. Published by St Lucie Press, Delray Beach, Florida. Pp. 325–331.
- Carvalho, C. R. L., Rossetto, C. J., Mantovani, D. M. B., Morgano, M. A., de Castro, J. V. & Bortoletto, N., 2004. Avaliação de cultivares de mangueira selecionadas pelo Instituto Agronômico de Campinas comparadas a outras de importância comercial. *Revista Brasileira de Fruticultura* 26: 264–71.
- Cazorla, F.M., Tores, J.A., Olalla, L., Perez-Garcia, A., Farre, J.M. & de Vicente, A., 1998. Bacterial apical necrosis of mango in southern Spain: a disease caused by *Pseudomonas syringae* pv. *syringae*. *Phytopathology* 8: 614–620.
- Cazorla, F. M., Arrebola, E., Olea, F., Velasco, L., Hermoso, J. M., Perez-Garcia, A., Tores, J. A., Farre, J. M. & de Vicente, A. 2006. Field evaluation of treatments for the control of the bacterial apical necrosis of mango (*Mangifera indica*) caused by *Pseudomonas syringae* pv. *syringae*. *European Journal of Plant Pathology* 116: 279–288.
- Chadha K.L. 1989. World mango industry. *Acta Horticulturae* 231: 3-11.
- Chavan, R.A., Deshmukh, V.D., Tawade, S.V. & Deshmukh, J.D., 2009. Efficacy of fungicides for managing powdery mildew of mango. *International Journal of Plant Protection* 2: 71-72.
- Cho, H.K. & Cho, J.Y., 2010. Cho Han-Kyu Natural Farming. Cho Global Natural Farming, Seoul, Korea.
- Colangelo, M. A., Caruso, M. C., Favati, F., Scarpa, T., Condelli, N. & Galgano, F., 2015. Electrolysed water in the food industry as supporting of environmental sustainability. In: "The Sustainability of Agro-Food and Natural Resource Systems in the Mediterranean Basin", edited by Vastola, A. Published by Springer International Publishing. Pp. 385–397.
- Colosimo, E. A., Chalita, L. V. A. S. & Demetrio, C. G. B., 2000. Tests of proportional hazards and proportional odds models for grouped survival data. *Biometrics* 56: 1233–1240.
- Cook, A.A., 1975. *Diseases of Tropical and Subtropical Fruits and Nuts*. Hafner Press, New York.
- Cooke, A. W., 2007. Phytophthora postharvest fruit rot in mango. Advisory poster produced by the Queensland Government Department of Primary Industries and Fisheries. Indoorpilly, Australia.
- Cooke, A.W., van der Kruyssen, A. & Johnson, G.I., 1998. The effect of commercial pruning on colonization of mango by endophytic *Dothiorella* species. In: "Post-harvest Handling of Tropical Fruit", edited by Champ, B.R., Highley, E. and Johnson, G.I. Australian Centre for International Agricultural Research (ACIAR), Canberra. Pp. 128–130.
- Coto, D., Saunders, J.L., Vargas, C.L. & King, A.B.S., 1995. Plagas invertebradas de cultivos tropicales con énfasis en América central. *Centro Agronómico Tropical de Investigación y Enseñanza. Serie Técnica, Manual Técnico* 12.
- Crespo, M., Cazorla, F. M., Hermoso, J. M., Guirado, E., Maymon, M., Tores, J. A., Freeman, S. & de Vicente, A., 2012. First report of mango malformation disease caused by *Fusarium mangiferae* in Spain. *Plant Disease* 96: 286–287.

- Cronje, C., Wehnwe, F. C. & Kotze, J. M. 1990. *Alternaria alternata* as a lesion pathogen of mango inflorescences in South Africa. *Phytophylactica* 22: 117–118.
- Crookes, C. A. & Rijkenberg, F. H. J., 1985. A literature review of the distribution, symptomatology, cause and control of mango blossom malformation. *South African Mango Growers Association Research Report* 5: 15–24.
- Cunningham, C., 1984. Mango insect pest. In: “Australian Mango Research Workshop”. 26–30 November 1984. CSIRO, Melbourne. Pp. 211-224.
- Dag, A., Eisenstein, D. & Gazit, S., 2001. Effects of three fungicides used to control powdery mildew in mango on pollen germination and pollen-tube growth. In: *Proceedings of the Interamerican Society for Tropical Horticulture*, vol. 43: 123-126.
- Daneel, M. & Joubert, P., 2009. Biological control of the mango scale *Aulacaspis tubercularis* Newstead (Coccidae: Diaspididae) by a parasitoid *Aphytis chionaspis* Ren (Hymenoptera: Aphelinidae). *Acta Horticulturae*, 820: 567–74.
- Darvas, J.M., 1987. Control of mango blossom malformation with trunk injection. *South African Mango Grower's Association Yearbook* 7: 21-24.
- Darvas, J.M., 1993. *Dothiorella dominicana*, an important mango pathogen in South Africa. *Acta Horticulturae* 341: 321–328.
- Das Gupta, S.N. & Zacchariah, H.T., 1945. Studies on the diseases of *Mangifera indica*. Part V. On the die-back of mango trees. *Journal of the Indian Botanical Society* 24, 101–108. (Abstract in *Review of Applied Mycology*, 25: 267–268.
- De, K. & Pande, Y.D., 1987. Evaluation of certain non-insecticidal methods of reducing infestation of the mango nut weevil, *Sternochetus gravis* (F.) in India. *Tropical Pest Management* 33: 27–28.
- De Faveri, S., 2018. Integrated pest management and biological pest control strategies in mango cultivation. In: “Pests and Their Management”, edited by Omkar. Published by Springer Nature Signature. Pp. 511-541. <http://dx.doi.org/10.19103/AS.2017.0026.21>.
- De Jesus, LRA & Cortez F., 1998. Field infestation and chemical control of the mango pulp weevil, *Sternochetus frigidus* (Fabr) in Palawan. *Philipp Agric* 81(3 & 4): 131-140.
- De la Rosa, W., Lopez, F.L. & Liedo, P., 2002. *Beauveria bassiana* as a pathogen of the Mexican fruit fly (Diptera: Tephritidae) under laboratory conditions. *Journal of Economic Entomology* 95: 36–43.
- De Laroussilhe, F., 1980. Le Manguier. G.P. Masoneuve & Larose, Paris.
- De Macedo, F.P., 1998. Efeitos da aplicacao de acido giberelico em manga (*Mangifera indica* L.) cultivar Tommy Atkins: aspectos horticulturais e entomologicos. Tesis de doctorado, Universidade de Sao Paulo, Brazil, 120 pp.
- Desai, S.A., 1998. Tridemefon in the control of powdery mildew of mango. *Karnataka Journal of Agricultural Science* 11: 244-245.
- Devi Thangam, S., Verghese, A., Dinesh, M.R., Vasugi, C. & Kamala Jayanthi, P.D., 2013. Germplasm evaluation of mango for preference of the mango hopper, *Idioscopus nitidulus* (Walker) (Hemiptera: Cicadellidae): The first step in understanding the host plant resistance. *Pest Manag Hort Ecosyst* 19: 10–16.
- Dias, N.P., Zotti, M.J., Montoya, P., Carvalho, I.R. & Nava, D.E., 2018. Fruit fly management research: a systematic review of monitoring and control tactics in the world. *Crop Protection* 112: 187-200.

- Díaz-Fleischer, F. & Aluja, M., 2003. Clutch size in frugivorous insects as a function of host firmness: the case of the tephritid fly *Anastrepha ludens*. *Ecological Entomology* 28: 268–277.
- Díaz-Fleischer, F., Toledo, J., Enkerlin, W. & Fernandez, J., 1996. Cyromazine: effects on three species of *Anastrepha*. In “Fruit Fly Pests: a World Assessment of their Biology and Management”, edited by McPherson, B. & Steck, G. Published by St Lucie Press, Delray Beach, Florida. Pp. 336–337.
- Díaz-Fleischer, F., Pérez-Staples, D., Cabrera-Mireles, H., Montoya, P. & Liedo, P., 2017. Novel insecticides and bait stations for the control of *Anastrepha* fruit flies in mango orchards. *J. Pest. Sci.* 90, 865–872.
- Dimbi, S., Maniania, N.K., Lux, S., Ekesi, S. & Mueke, J.K., 2003. Pathogenicity of *Metarhizium anisopliae* (Metsch.) Sorokin and *Beauveria bassiana* (Balsamo) Vuillemin, to three adult fruit fly species: *Ceratitis capitata* (Wiedemann), *C. rosa* var. *fasciventris* Karsch and *C. cosyra* (Walker) (Diptera: Tephritidae). *Mycopathologia* 156: 375–382.
- Dodd, J.C., Estrada, A.B., Matcham, J., Jeffries, P. & Jeger, M.J., 1991a. The effect of climatic factors on *Colletotrichum gloeosporioides*, the causal agent of mango anthracnose, in the Philippines. *Plant Pathology* 40: 568–575.
- Dodd, J.C., Bugante, R., Koomen, I., Jeffries, P. & Jeger, M.J., 1991b. Pre- and post-harvest control of mango anthracnose in the Philippines. *Plant Pathology* 40: 576–583.
- Dodd, J.C., Prusky, D. & Jeffries, P., 1997. Fruit diseases. In: “The Mango: Botany, Production and Uses”, edited by R.E. Litz. Published by CAB International, Wallingford, UK. Pp. 257–280.
- Domsch, K.H., Gams, W. & Anderson, T.H., 1980. Compendium of Soil Fungi. Academic Press, London.
- Dos Santos, M.G., Gonçalves, A.C. Jr, Junior, D.L.B., Schwantes, D., Campagnolo, M.A., Coelho, G.F., Parizotto, A.A, Zimmermann, J. & Vargas de Toledo, M., 2016. Utilização de medicamentos homeopáticos para remediação de solos contaminados por metais tóxicos. *Journal of Agronomic Sciences*, 5: 26-45.
- Droby, S., Wisniewski, M., Teixido, N., Spadaro, D. and Jijakli, M. H., 2016. The science, development, and commercialization of postharvest biocontrol products. *Postharvest Biology and Technology* 122: 22–29.
- Epsky, N., Hendrichs, J., Katsoyannos, B., Vasquez, A., Ros, J., Zumreolu, A., Bakri, A., Seewooruthum, S. & Heath, R., 1999. Field evaluation of female-targeted trapping systems for *Ceratitis capitata* (Diptera: Tephritidae) in seven countries. *Journal of Economic Entomology* 92: 156–164.
- Eshel, D., Miara, I., Ailing, T., Dinoor, A. & Prusky, D., 2002. pH regulates endoglucanase expression and virulence of *Alternaria alternata* in persimmon fruits. *Molecular Plant Microbe Interactions* 15: 774–779.
- Estrada, A.B., Jeffries, P. & Dodd, J.C., 1996. Field evaluation of a predictive model to control anthracnose disease of mango in the Philippines. *Plant Pathology* 45: 294–301.
- Fallik, E., 2004. Prestorage hot water treatments (immersion, rinsing and brushing). *Postharvest Biology and Technology* 32: 125–134.
- FAO/IAEA, 2019. Use of Entomopathogenic Fungi for Fruit Fly Control in Area-Wide SIT Programmes, edited by Villaseñor, A. Flores, S., Campos, S.E., Toledo, J.P., Montoya, P., Liedo, P. and Enkerlin, W. Published by Food and Agriculture Organization of the United Nations/ International Atomic Energy Agency. Vienna, Austria. 43pp.
- Fasih, M. & Srivastava, R.P., 1990. Parasites and predators of insect pests of mango. *Integrated Pest Control* 32: 39–41.

- FDA (Food and Drugs Administration), 2020. <https://www.fda.gov/drugs/information-drug-class/homeopathic-products>
- Fitzell, R. D., Peak, C. M. & Darnell, R. E., 1984. A model for estimating infection levels of anthracnose disease of mango. *Annals of Applied Biology* 104: 451–458.
- Flechtmann, C. H. W., Kimati, H., Medcalf, J. C. & Ferre, J., 1973. Preliminary observations on malformation in mango (*Mangifera indica* L.) inflorescences and the fungus, insects and mite found on them. *Review of Plant Pathology* 52: 162–163.
- Flores, S., Montoya, P., Toledo, J. & Enkerlin, W., 2014. Estimation of populations and sterility induction in *Anastrepha ludens* (Diptera: Tephritidae) fruit flies. *J. Econ. Entomol.* 107: 1502–1507.
- Flores, S., Gómez-Escobar, E., Liedo, P., Toledo, J. & Montoya, P., 2017. Density estimation and optimal sterile-to-wild ratio to induce sterility in *Anastrepha obliqua* populations. *Entomol. Exp. Appl.* 164: 284–290.
- Follett, P. A., 2001. Irradiation as a quarantine treatment for mango seed weevil (Coleoptera: Curculionidae). *Proc. Hawaii Entomol. Soc.* 35: 95–100.
- Follett, P. & Neven, L.G., 2006. Current trends in quarantine entomology. *Annual Review of Entomology* 51: 359–385.
- Freeman, S., Otero-Colina, G., Rodriguez-Alvarado, G., Fernandez-Pavia, S., Maymon, M., Ploetz, R.C., Aoki, T. & O'Donnell, K., 2014a. First report of mango malformation disease caused by *Fusarium pseudocircinatum* in Mexico. *Plant Disease* 8: 1583.
- Freeman, S., Shtienberg, D., Maymon, M., Levin, A. G. & Ploetz, R. C., 2014b. New insights into mango malformation disease epidemiology lead to a new integrated management strategy for subtropical environments. *Plant Disease* 98: 1456–1466.
- Gagné, R.J., & Medina, C., 2004. A new species of Procontarinia (Diptera: Cecidomyiidae), an important new pest in the Philippines. *Proceedings of the Entomological Society of Washington* 106: 19–25.
- García-Medel, D., Sivinski, J., Díaz-Fleischer, F., Ramírez-Romero, R. & Aluja, M., 2007. Foraging behavior by six fruit fly parasitoids (Hymenoptera: Braconidae) released as single- or multiple-species cohorts in field cages: influence of fruit location and host density. *Biological Control* 43: 12–22.
- García-Ramírez, M.J., Cibraian-Tovat, J., Arzufi-Barrera, R., López-Collado, J. & Soto-Hernández, M., 2004. Preferencia de *Anastrepha ludens* (Loew) (Diptera: Tephritidae) por volátiles de frutos verdes o amarillos de mango y naranja. *Agrociencia* 38: 423–430.
- Giardini-Bonfim, F.P., Dias-Casali, V.W. & Ronie-Martins, E., 2012 Germinação e vigor de sementes de tomate (*Lycopersicon esculentum* Mill), peletizadas com preparados homeopáticos de *Natrum muriaticum*, submetidas a estresse salino. *Enciclopédia Biosfera*, Centro Científico Conhecer-Goiânia, 8 (14): 625-633.
- Godase, S.K., Bhole, S.R., Shivpuje, P.R. & Patil, B.P., 2004. Assessment of yield loss in mango (*Mangifera indica*) due to mango hopper (*Idioscopus niveoparvus*) (Homoptera: Cicadellidae). *Indian Journal of Agricultural Sciences* 74: 370–372.
- Golez, H.G., 1991. Bionomics and control of the mango seed borer, *Noorda albizonalis* Hampson (Pyralidae: Lepidoptera). *Acta Horticulturae* 291: 418–424.
- Gagnevin, L. & Pruvost, O., 2001. Epidemiology and control of mango bacterial black spot. *Plant Disease* 85: 928–935.

- Govender, V., Korsten, L. & Sivakumar, D., 2005. Semi-commercial evaluation of *Bacillus licheniformis* to control mango postharvest diseases in South Africa. *Postharvest Biology and Technology*, 48: 254–258.
- Guentzel, J. L., Lam, K. L., Callan, M. A., Emmons, S. A. & Dunham, V. L., 2010. Postharvest management of gray mold and brown rot on surfaces of peaches and grapes using electrolyzed oxidizing water. *International Journal of Food Microbiology* 143: 54–60.
- Gupta, J.H. & Yadav, A.S., 1984. Chemical control of powdery mildew of mango. *Indian Journal of Mycology and Plant Pathology* 14: 297-298.
- Haji, F. N. P., Barbosa, F. R., Lopes, P. C. R., Moreira, A. N., de Alencar, J. A. & Ferreira, R. C. F., 2004. Monitoring mango pests within an integrated production program in Brazil. *Acta Horticulturae* 645: 163–165.
- Hallman, G.J. & Sharp, J.L., 1990. Mortality of Caribbean fruit fly (Diptera: Tephritidae) larvae infesting mangoes subjected to hot-water treatment, then immersion cooling. *Journal of Economic Entomology* 83: 2320–2323.
- Hansen, J.D., 1993. Dynamics and control of the mango seed weevil. *Acta Horticulturae* 341: 415–420.
- Hansen, J.D. & Armstrong, J.W., 1990. The failure of field sanitation to reduce infestation by the mango nut weevil, *Cryptorhynchus mangiferae* (F.) (Coleoptera: Curculionidae). *Tropical Pest Management* 36: 359–361.
- Heath, R.R., Epsky, N.D., Guzman, A., Dueben, B.D., Manukian, A. & Meyer, W.L., 1995. Development of a dry plastic insect trap with food-based synthetic attractant for the Mediterranean and Mexican fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology* 88: 1307–1315.
- Heath, R.R., Epsky, N.D., Dueben, B.D., Rizo, J. & Jeronimo, F., 1997. Adding methyl substitute ammonia derivatives to food-based synthetic attractant on capture of the Mediterranean and Mexican fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology* 90: 1584–1589.
- Hennesey, M. & Schnell, R.J., 2001. Resistance of immature mango fruits to Caribbean fruit fly (Diptera: Tephritidae). *Florida Entomologist* 84: 318–319.
- Hernandez-Sanchez, G., Sanz-Barbosa, I., Casana-Giner, V. & Primo-Yufera, E., 2001. Attractiveness for *Ceratitidis capitata* (Wiedemann) (Dipt., Tephritidae) of mango (*Mangifera indica*, cv. Tommy Atkins) airborne terpenes. *Journal of Applied Entomology* 125: 189–192.
- Herrera, A.J. & Vinas, L.E., 1977. Fruit flies (Diptera: Tephritidae) on mangoes in Chulucanas, Piura. *Revista Peruana de Entomologia* 20: 107–114.
- Hingorani, M.K., Sharma, O.P. & Sohi, H.S., 1960. Studies on blight disease of mango caused by *Macrophoma mangiferae*. *Indian Phytopathology* 13: 137–143.
- Hoffman, H., 1860. Untersuchungen über die Keimung der Pilzsporen. *Jb. Wiss. Bot.*, 2: 267-337.
- Hofman, P.J., Smith, L.G., Joyce, D.C., Johnson, G.I. & Meiburg, G.F., 1997. Bagging of mango (*Mangifera indica* cv. 'Keitt') fruit influences fruit quality and mineral composition. *Postharvest Biology and Technology* 12: 83–91.
- Holliday, P., 1980. *Fungus Diseases of Tropical Crops*. Cambridge University Press: Cambridge.
- Honger, J.O., Offei, S.K., Oduro, K.A., Odamtten, G.T., & Nyaku, S.T., 2014. Identification and species status of the mango biotype of *Colletotrichum gloeosporioides* in Ghana. *European Journal of Plant Pathology* 140 (3): 455–467.

- Hsu, J.-C. & Feng, H.-T. (2006) Development of resistance to Sinosad in oriental fruit fly (Diptera: Tephritidae) in laboratory selection and cross-resistance. *Journal of Economic Entomology* 99: 931–936.
- Huete, M. & Arias, S., 2007. Manual para la Producción de Mango. USAID-RED. Proyecto de Diversificación Económica Rural.
- Hughes, S.J., 1980. New Zealand Fungi 29. *Rhinocladium* Sacc. et March. *New Zealand Journal of Botany* 18: 163–172.
- Hussain, S., Masud, T. & Ahad, K., 2002. Determination of pesticides residues in selected varieties of mango. *Pakistan Journal of Nutrition* 1(1): 41-42.
- IAEA, 2003. Trapping Guidelines for Area-wide Fruit Fly Programmes. International Atomic Energy Agency, Vienna.
- ICAR, 2014. Management of mango trunk borer (*Batocera rufobaculata*). <http://www.cish.res.in/Technologies/Management%20of%20Stem%20borer%20final%20%20English.pdf>.
- Ihsan, J., Ahmad, I., Sajid, M.N., Muhammad, F. & Saleem, A., 1999. Incidence of powdery mildew of mango in the Punjab and evaluation of protective and curative fungicides for the control of disease. *Pakistan Journal of Phytopathology* 11: 67-69.
- Iqbal, Z, Akhtar, N., Ghazanfar, M.U., Shehzad, S.M., Ahmad, S., Asif, M., Yasin, M., Pervez, M.A., Dasti, A.A. & Saleem, A., 2011. Management of mango malformation through physical alteration and chemical spray. *African Journal of Agricultural Research* 6 (7): 1897-1901.
- Islam, M.S. & Elegio, D.T., 1997. Effect of time and frequency of insecticide spraying on the control of mango leafhoppers (*Idioscopus* spp.). *Bangladesh Journal of Entomology* 7: 93–99.
- Istianto, M. & Soemargono, A., 2015. The effect of citronella essential oil on controlling the mango red-banded caterpillar, *Noorda albizonalis* Hampson (Lepidoptera: Pyralidae). *Jordan Journal of Biological Sciences* 8(2): 77-80.
- Iyer, C. P. A., 2004. Growing mango under organic system. *Acta Horticulturae* 645: 71–84.
- Iwahashi, O., 1984. The control project of the oriental fruit fly in Okinawa. *Chinese Journal of Entomology* 4: 107–120. (in Chinese)
- Janisiewicz, W.J. & Conway, W.S., 2010. Combining biological control with physical and chemical treatments to control fruit decay after harvest. *Stewart Postharvest Review* 6 (1): 1–16.
- Janisiewicz, W.J., Jurick Li, W.M., Vico I., Peter K.A. & Buyer J.S., 2013. Culturable bacteria from plum fruit surfaces and their potential for controlling brown rot after harvest. *Post-harvest Biology and Technology* 76: 145–151.
- Jayanthi, P.D.K., Verghese, A., Shashank. P.R. & Kempraj. V., 2014a. Spread of indigenous restricted fruit borer, *Citripestis eutrappera* (Meyrick) (Pyralidae: Lepidoptera) in mango-time for domestic quarantine regulatory reforms. *Pest Manag Hort Ecosyst* 20(2): 227–230.
- Jayanthi, P.D.K., Verghese, A., Arthikruba, A., Sowmya, B.R. & Bhatt, R.M., 2014b. Emerging pests of mango under changing climate scenario, Technical bulletin no. 18. Indian Institute of Horticultural Research, Bengaluru.
- Jeffries, P., Dodd, J. C., Jeger, M. J. & Plumbley, R. A., 1990. The biology and control of *Colletotrichum* species on tropical fruit crops. *Plant Pathology* 39: 343–366.

- Jeffries, P. & Koomen, I., 1992. Strategies and prospects for biological control of diseases caused by *Colletotrichum*. In: "Colletotrichum: Biology, Pathology and Control", edited by Bailey, J.A. and Jeger, M.J. Published by CAB International, Wallingford, UK. Pp. 337–357.
- Jhala, R.C., Shah, A.H., Patel, Z.P. & Patel, R.L., 1989. Studies on population dynamics of mango hopper and scope of spraying in integrated pest management programme. *Acta Horti* 231: 597–601.
- Jiron, L.F., 1995. Opciones al uso de insecticidas en mango. In: "Opciones al uso Unilateral de Plaguicidas en Costa Rica. Vol. 2", edited by Garcia, J., Fuentes, G. and Monge, J. Published by Editorial Euned, San Jose, Costa Rica. Pp. 129–155.
- Jiron, L.F. & Soto-Manitiu, J., 1987. Las Moscas de las frutas (Diptera: Tephritidae) en Costa Rica: situacion actual, analisis y comentario. *Agronomia Costarricense* 11: 255–261.
- Joel, D.M., 1980. Resin ducts in the mango fruit: a defense system. *Journal of Experimental Botany* 30: 1707–1718.
- Johnson, G.I., 1994a. Powdery mildew. In "Compendium of Tropical Fruit Diseases", edited by Ploetz, R.C., Zentmyer, G.A., Nishijima, W., Rohrbach, K. and Ohr, H.D. Published by American Phytopathological Society (APS) Press, St Paul, Minnesota, USA. Pp. 38–39.
- Johnson, G.I., 1994b. Stem-end rot. In "Compendium of Tropical Fruit Diseases", edited by Ploetz, R.C., Zentmyer, G.A., Nishijima, W., Rohrbach, K. and Ohr, H.D. Published by American Phytopathological Society (APS) Press, St Paul, Minnesota, USA. Pp. 39–41.
- Johnson, G.I., Muirhead, I.F., Mayers, P. & Cooke, T., 1989. Diseases. In: "Mango Pests and Disorders", edited by Ridgeway, R. Department of Primary Industries, Brisbane, Queensland. Pp. 1–9.
- Johnson, G.I., Boag, T.S., Cooke, A.W., Izard, M., Panitz, M. & Sangchote, S., 1990. Interaction of post-harvest disease control treatments and gamma irradiation in mangoes. *Annals of Applied Biology*, 116 (2): 245-257.
- Johnson, G.I., Cooke, A.W., Mead, A.J. & Wells, L.A., 1991. Stem end rot of mango in Australia: causes and control. *Acta Horticulturae* 219: 288–295.
- Johnson, G.I., Mead, A.J., Cooke, A.W. & Dean, J.R., 1992. Mango stem end rot pathogens – fruit infection by endophytic colonization of the infl orescence and pedicel. *Annals of Applied Biology* 120: 225–234.
- Johnson, E.M., Sutton, T.B. & Hodges, C.S., 1997. Etiology of apple sooty blotch disease in North Carolina. *Phytopathology* 87, 88–95.
- Jones, R.W. & Prusky, D., 2001. Expression of an antifungal peptide in *Saccharomyces*: a new approach for biocontrol of the postharvest disease caused by *Colletotrichum coccodes*. *Phytopathology* 92: 33–37.
- Joubert, M. H., 1991. Implications of epidemiological studies on strategies for control of powdery mildew and anthracnose. *Yearbook South African Mango Growers Association*, 11: 26-28.
- Joubert, T.H. & Labuschagne, T.I., 1995. Alternative measures for controlling mango seed weevil, *Sternochetus mangiferae* (F). *Yearbook South African Mangos' Grower Association*, 15: 94-96.
- Joubert, P., Daneel, M. & Grove, T., 2000. Progress towards integrated pest management (IPM) on mangoes in South Africa. *Acta Horticulturae*, 509: 811–17.
- Junqueira, N. T. V., Pinto, A. C.de Q., da Cunha, M. M. & Ramos, V. H. V., 2002. Controle das doenças da mangueira. In: "Controle de Doenças de Plantas Fruteiras. Vol. 1", edited by Zambolin, L., do Vale, F. X. R., Monteiro, A. J. A. and Costa, H. Universidade Federal de Vicosa, Vicosa. Pp. 323-403.

- Kadman, A. & Gazit, S., 1984. The problem of iron deficiency in mango trees and experiments to cure it in Israel. *Journal of Plant Nutrition* 7: 282–90.
- Kamil, F.H., Saeed, E.E., El-Tarabily, K.A. & AbuQamar, S.F., 2018. Biological control of mango dieback disease caused by *Lasiodiplodia theobromae* using streptomycete and non-streptomycete actinobacteria in the United Arab Emirates. *Front. Microbiol.* 9: 829. doi: 10.3389/fmicb.2018.00829.
- Katoch, P., Katoch, A. & Dangi, P., 2019. An overview on mango malformation and the potential approaches to their management. *Journal of Pharmacognosy and Phytochemistry* 8 (4): 621-626.
- Kaur, L. Gupta, B., Sharma, I.M., & Joshi, A.K., 2018. Eco-friendly management of powdery mildew of mango through biocontrol agents. *International Journal of Current Microbiology and Applied Sciences* 7 (10): 392-396.
- Kawate, M., 1993. Pesticides registered for mango. In: “Proceedings: Conference on Mango in Hawaii”, edited by Chia, C.L., Evans, D.O. University of Hawaii at Manoa. Pp. 25-27.
- Keliikuli, A., Smith, K., Li, Y. & Lee, C.N., 2019. Natural Farming: The Development of Indigenous Microorganisms Using Korean Natural Farming Methods. *Sustainable Agriculture*, Dec. 2019. Pp 9.
- Kehri, K.H. & Chandra, S., 1986. Control of *Botryodiplodia* rot of guava with a homeopathic drug. *Natl Acad Sci Lett*, 9: 301–302.
- Khanzada, A.G. & Naqvi, K.M., 1985. The optimum time for the control of mango hopper *Idioscopus* spp. (Hemiptera: Cicadellidae). *Proceedings of the Entomological Society of Karachi* 14/15: 149–156.
- Khanna, K.K. & Chandra, S., 1978. A homoeopathic drug controls mango fruit rot caused by *Pestalotia mangiferae* Henn. *Experientia* 34: 1167–1168.
- King, J.R. & Hennessey, M.K., 1996. Spinosad bait for the Caribbean fruit fly (Diptera: Tephritidae). *Florida Entomologist* 79: 526–531.
- Kishun, R., 1994. Evaluation of phylloplane micro-organisms from mango against *Xanthomonas campestris* pv. *mangiferaeindicae*. *Indian Phytopathology* 47: 313.
- Kok I. B., 1979. Control of the mango seed weevil by trapping and irradiation. (Beheer van die mangosnuitkewer met klem op bestraling.). *Citrus and Subtropical Fruit Journal*. 14-16.
- Kolesik, P., Rice, A., Bellis, G. & Winthersohn, M., 2009. Procontarinia pustulata, a new gall midge species (Diptera: Cecidomyiidae) feeding on mango, *Mangifera indica* (Anacardiaceae), in northern Australia and Papua New Guinea. *Australian Journal of Entomology* 48(4): 310 – 316.
- Korsten, L. 2004. Biological control in Africa: Can it provide a sustainable solution for control of fruit diseases?. *South African Journal of Botany* 70 (1): 128–39.
- Korsten, L., 2006. Advances in control of postharvest diseases in tropical fresh produce. *International Journal of Postharvest Technology and Innovation* 1: 48–61.
- Kudagamage, C., de Z. Rajapakse, H.L., Rajapakse, R.H. & Ratnasekara, D., 2001. The population dynamics and insecticidal control of three leaf hoppers, *Amrytodus brevistilus* Leth., *Idioscopus niveosparus* Leth., and *Idioscopus clipealis* Leth. (Homoptera: Cicadellidae) in mango. *Journal of Entomological Research* 25: 121–125.
- Kueprakone, U., Saengkong, S., Pienpuck, K. & Choobumroong, W., 1986. *Phytophthora palmivora* (Butl.) Butl., the causal organism of black rot of mango seedlings. *Thai Agricultural Research Journal* 4: 67–73.

- Kumar, J. & Beniwal, S.P.S., 1991. Mango malformation. In "Plant Diseases of International Importance. Vol. III: Diseases of Fruit Crops", edited by Kumar, J., Chaube, H.S., Singh, U.S. and Mukhopadhyay, A.N. Published by Prentice Hall, Englewood Cliffs, New Jersey, USA. Pp. 357–393.
- Kumar, S., Rai, A., Patel, C. & Bhatl, R., 1993. Studies on spatial distribution of *Amritodes atkinsoni* (Leth.) infesting mango in south Gujarat, India. (Abstract). *IV International Mango Symposium*, p. 122.
- Kumar, P., Misra, A.K., Pandey, B.K., Misra, S.P. & Modi, D.R., 2009. Eco-friendly management of mango malformation (*Fusarium moniliforme* var. *subglutinans* = *Fusarium subglutinans*) through certain plant leaf extracts. *Journal of Eco-friendly Agriculture* 4 (1): 61-64.
- Kumar, P., Misra, A.J., Raj Modi, D.R. & Gupta, V.K. 2012. Biocontrol potential of Trichoderma species against mango malformation pathogens. *Archives of Phytopathology and Plant Protection*, 45 (10): 1237-1245.
- Kvas, M., Steenkamp, E. T., Al Adawi, A. O., Deadman, M. L., Al Jahwari, A. A., Marasas, W. F. O., Wingfield, B. D., Ploetz, R. C. & Wingfield, M. J., 2008. *Fusarium mangiferae* associated with mango malformation in the Sultanate of Oman. *European Journal of Plant Pathology* 121: 195–199.
- Lasa, R., Ortega, R. & Rull, J. 2013. Towards development of a mass trapping device for Mexican fruit fly *Anastrepha ludens* (Diptera: Tephritidae) control. *Florida Entomologist* 96: 1135–1142.
- Le Lagadec, M. D., 2004. The control of mango scale in commercial orchards through the use of the predatory beetle *Cybocephalus binotatus*. *Acta Horticulturae* 645: 509–15.
- Lezama-Gutierrez, R., Trujillo, A., Molina-Ochoa, J., Rebolledo-Dominguez, O., Pescador, A., Lopez-Edwards, M. & Aluja, M., 2000. Virulence of *Metarrizium anisopliae* (Deuteromycotina: Hypomycetes) on *Anastrepha ludens* (Diptera: Tephritidae): laboratory and field trials. *Journal of Economic Entomology* 93: 1080–1084.
- Lezama-Gutiérrez, R., Molina-Ochoa, J., Pescador-Rubio, A., Galindo-Velasco, E., Ángel-Sahagún, C.A., Michel-Aceves, A.C. & González-Reyes, E., 2006. Efficacy of steinernematid nematodes (Rhabditida: Steinernematidae) on the suppression of *Anastrepha ludens* (Diptera: Tephritidae) larvae in soil of differing textures: laboratory and field trials. *J. Agric. Urban Entomol.* 23: 41–49.
- Liburd, O.E., Holler, T.C. & Moses, A.L., 2004. Toxicity of Imidacloprid-treated spheres to Caribbean fruit fly, *Anastrepha suspensa* (Diptera: Tephritidae) and its parasitoid *Diachasmimorpha longicaudata* (Hymenoptera: Braconidae) in the laboratory. *Journal of Economic Entomology* 97: 525–529.
- Liew, E. C. Y., Laurence, M. H., Pearce, C., Shivas, R. G., Johnson, G. I., Tan, Y. P., Edwards, J., Perry, S., Cooke, A. W. & Summerell, B. A., 2016. Review of *Fusarium* species isolated in association with mango malformation in Australia. *Australasian Plant Pathology* 45: 547–559.
- Lim, T.C. & Khoo, K.C., 1985. Diseases and disorders of mango in Malaysia. Tropical Press: Kuala Lumpur, Malaysia.
- Lima, C. S., Pfenning, L. H., Costa, S. S., Abreu, L. M. & Leslie, J. F., 2012. *Fusarium tuiense* sp. nov., a member of the *Gibberella fujikuroi* complex that causes mango malformation in Brazil. *Mycologia* 104: 1408–1419.
- Lima, N.B., Batista, M.V.A., De Morais, M.A., Jr., Barbosa, M.A.G., Michereff, S.J., Hyde, K.D., & Câmara, M.P.S., 2013. Five *Colletotrichum* species are responsible for mango anthracnose in northeastern Brazil. *Fungal Diversity* 61: 75–88
- Lima, G. P. P., Machado, T. M., de Oliveira, L. M., da Silva Borges, L., de Albuquerque Pedrosa, V., Vanzani, P. & Vianello, F., 2014. Ozonated water and chlorine effects on the antioxidant properties of organic and conventional broccoli during postharvest. *Scientific Agriculture* 71: 151–156.

- Lindgren, J. & Vail, P., 1986. Susceptibility of Mediterranean fruit fly, melon fly and oriental fruit fly (Diptera: Tephritidae) to the entomogenous nematode *Steinernema feltiae* in laboratory tests. *Environmental Entomology* 4: 465–468.
- Lonsdale, J. H. & Kotze, J. M., 1993. Chemical control of mango blossom diseases and the effect on fruit set and yield. *Plant Disease* 77: 558–562.
- Lopez, F., Chambers, D., Sanchez, M. & Kamasaki, H., 1969. Control of the Mexican fruit fly by bait sprays concentrated at discrete locations. *Journal of Economic Entomology* 62: 1255–1257.
- Lopez, M., Aluja, M. & Sivinski, J., 1999. Hymenopterous larval-pupal and pupal parasitoids of *Anastrepha* flies (Diptera: Tephritidae) in Mexico. *Biological Control* 15: 119–129.
- Lourd, M. & Keuli, S. D., 1975. Note sur un chancre a *Phytophthora* du manguier en Cote d'Ivoire. *Fruits* 30: 541–544.
- Louw, C.E., 2009. Biology and control of the mango seed weevil in South Africa. Department of Zoology and Entomology. Faculty of Natural and Agricultural Sciences University of the Free State of Bloemfontein.
- Louw, C.E., 2013. The mango seed weevil: *Sternochetus mangiferae* (Fabricius) (Coleoptera: Curculionidae); understanding the pest in order to ensure effective control measures. *Acta Horti* 992: 441-457.
- Love, K., Bowen, R. & Paull, R., 2003. Increasing marketable production of exotic tropical fruit with protective covering. *Western Region, Sustainable Agriculture Research and Education, Project Report, Farmer/Rancher, Sustainable Agriculture Research and Education, Utah State University, Captain Cook, Hawaii, USA.*
- Luo, S., Wan, B., Feng, S. & Shao, Y., 2015. Biocontrol of postharvest anthracnose of mango fruit with *debaryomyces nepalensis* and effects on storage quality and postharvest physiology. *Journal of Food Science* 80: 2555–2563.
- Lurie, S., 1998. Postharvest heat treatments. *Postharvest Biology and Technology* 14: 257–69.
- Lux, S.A., Ekesi, S., Dimbi, S., Mohamed, S. & Billah, M., 2003. Mango-infesting fruit flies in Africa: perspectives and limitations of biological approaches to their management. In: “Biological Control in IPM Systems in Africa”, edited by Neuenschwander, P., Borgemeister, C. and Langewald, J. Published by CAB International, Wallingford, UK. Pp. 277–293.
- Malaguti, G. & de Reyes, C., 1964. A gall disease of cacao and mango in Venezuela caused by *Calonectria rigidiuscula*. *Phytopathology* 54: 499.
- Malo E.A. & Zapien, G.I., 1994. McPhail trap captures of *Anastrepha obliqua* and *Anastrepha ludens* (Diptera: Tephritidae) in relation to time of day. *Florida Entomologist* 77: 290–294.
- Mangan, R.L. & Hallman, G.H., 1998. Temperature treatments for quarantine security: new approaches for fresh commodities. In: “Temperature Sensitivity in Insects and Application in Integrated Pest Management”, edited by Hallman, G.H. and Denlinger, D.L. Published by Westview Press, Boulder, Colorado. Pp. 201–234.
- Mangan, R.L. & Moreno, D.S., 2007. Development of bait stations for fruit fly population suppression. *Journal of Economic Entomology* 100: 440–450.
- Mangan, R.L. & Sharp, J.L., 1994. Combination treatments. In: “Quarantine Treatments for Pests of Food Plants”, edited by Sharp, J.L. and Hallman, G.H. Published by Westview Press, Boulder, Colorado. Pp. 239–247.

- Mangan, R.L., Moreno, D.S., & Thompson, G.D., 2006. Bait dilution, spinosad concentration, and efficacy of GF-120 based fruit fly sprays. *Crop Protection* 25: 125–133.
- Manicom, B. Q. & Pruvost, O. P., 1994. Bacterial black spot. In “Compendium of Tropical Fruit Diseases”, edited by Ploetz, R. C., Zentmyer, G. A., Nishijima, W. T., Rohrbach, K. G. and Ohr, H. D. Published by APS Press, St Paul, USA. Pp. 41-42.
- Manoukis, N.C., Vargas, R.I., Carvalho, L., Fezza, T., Wilson, S., Collier, T. & Shelly, T.E., 2019. A field test on the effectiveness of male annihilation technique against *Bactrocera dorsalis* (Diptera: Tephritidae) at varying application densities. *PLOS ONE* 14(3): e0213337.
- Marasas, W. F. O., Ploetz, R. C., Wingfield, M. J., Wingfield, B. D. & Steenkamp, E. T., 2006. Mango malformation disease and the associated *Fusarium* species. *Phytopathology* 96: 667–72.
- Marlatt, R.B., Knight, R.J., Jr & Goldweber, S., 1970. *Verticillium* wilt of mango (*Mangifera indica*) in Florida. *Plant Disease Reporter* 54: 569–571.
- MARNDR (Ministry of Agriculture, Natural Resources, and Rural Development), 2014. Données d’identification des insectes capturés (2008–2014). Division of Plant Protection (DPV), Port-au-Prince, Haiti.
- Martínez-Ferrer, M.T., Campos, J.M. & Fibla, J.M., 2012. Field efficacy of *Ceratitis capitata* (Diptera: Tephritidae) mass trapping technique on clementine groves in Spain. *Journal of Applied Entomology* 136 (3): 161-240.
- Martorell, L.F., 1975. Annotated Food Plant Catalog of the Insects of Puerto Rico. Agricultural Experiment Station, University of Puerto Rico, Rio Piedras, Puerto Rico.
- Matheron, M. E. & Matejka, J. C., 1988. Phytophthora crown and root rot on nursery-grown mango trees delivered to Arizona. *Phytopathology* 78: 1572.
- Mazón-Suástegui, J.M., Ojeda-Silvera, C. M., García-Bernal, M., Avilés-Quevedo, M. A., Abasolo-Pacheco, F., Batista-Sánchez, D., Tovar-Ramírez, D., Arcos-Ortega, F., Murillo-Amador, B., Nieto-Garibay, A., Ferrer-Sánchez, Y., Morelos-Castro, R. M., Alvarado-Mendoza, A., Díaz-Díaz, M. & Bonilla-Montalvan, B., 2019. Agricultural Homoeopathy: A New Insight into Organics [Online First], IntechOpen, DOI: 10.5772/intechopen.84482.
- McGrath, M. & Andrews, J. 2005. What an epiphyte can teach us about foliar biocontrol: *Aureobasidium pullulans* on apple leaves as a model system. *Phytopathology* 95 (6): 568–568.
- McQuate, G.T., Peck, S.L., Barr, P.G. & Sylva, C.D., 2005. Comparative evaluation of spinosad and phloxine B as toxicants in protein baits for suppression of three fruit fly (Diptera: Tephritidae) species. *Journal of Economic Entomology* 98, 1170–1178.
- McSorley, R, Parrado, J. L. & Goldweber, S., 1980. Observations on a mango decline in South Florida. *Proceedings of the Florida State Horticultural Society* 93: 132–133.
- Medina, C.D.R., Ronquillo, S.J.Q., Mendiolo, M.S., Kolesik, P. & Newton I. (2017) Morphological and molecular characterization of the various species of *Procontarinia* (Diptera: Cecidomyiidae) attacking mango in the Philippines. *Acta Horticulturae* 1183: 287-290.
- Medina-Urrutia, V.M., Reyes-Hernández, J.E., Virgen-Calleros, G. & Pimienta-Barrios, E., 2018. Organic mango production. A review. In: “Achieving sustainable cultivation of mangoes”, edited by Galán-Saúco, V. and Lu, P. Published by Burleigh Dodds Science, Cambridge, UK. Pp. 229-268.
- Mertilus, F., Peña, J.E., Ring, D. & Schowalter, T., 2017. Inexpensive artisanal traps for mass trapping of fruit flies (Diptera: Tephritidae) in Haiti. *Florida Entomologist* 100(2): 390-395.

- Minas, I. S., Tanou, G., Belghazi, M., Job, D., Manganaris, G. A., Molassiotis, A. & Vasilakakis, M., 2012. Physiological and proteomic approaches to address the active role of ozone in kiwifruit post-harvest ripening. *Journal of Experimental Botany* 63: 2449–2464.
- Misra, A. K. & Prakash, O., 1992. Control of bacterial canker of mango by chemicals. *Pesticides* 18: 32–33.
- Misra, A., Pandey, D. & Singh, V.K., 2000. Mango malformation- An overview. In “Advances in Plant Diseases Mangement”, edited by Kumar, U., Kumar, K. and Srivastava, M. Published by Advance Publishing Concept. Pp. 185- 214.
- Modolon, T.A.; Boff, P., Boff, M.I.C. & Miquelluti, D.J., 2012. Homeopathic and high dilution preparations for pest management to tomato crop under organic production system. *Horticultura Brasileira*, 30: 51-57.
- Mohyuddin, A.I., 1981. A review of biological control in Pakistan. In: “Proceedings of the Second Pakistan Congress of Zoology”, Tandojam, Pakistan. Pp. 31–79.
- Mohyuddin, A.I. & Mahmood, R., 1993. Integrated control of mango pests in Pakistan. *Acta Horticulturae* 341: 467–483.
- Monaco, K. D. A., Costa, S. M., Minatel, L. O., Correa, C. R., Calero, F. A., Vianellod, F. & Lima, G. P. P., 2016. Influence of ozonated water sanitation on postharvest quality of conventionally and organically cultivated mangoes after postharvest storage. *Postharvest Biology and Technology* 120: 69–75.
- Montoya, P., Liedo, P., Benrey, B., Cancino, J., Barrera, J.F., Sivinski, J. & Aluja, M., 2000. Biological control of *Anastrepha* spp. (Diptera: Tephritidae) in mango orchards through augmentative releases of *Diachasmimorpha longicaudata* (Ashmead) (Hymenoptera: Braconidae). *Biological Control* 18: 216–224.
- Mossler M. A. & Nesheim, O.N., 2002. Florida Crop/Pest Management Profile: Mango. <http://edis.ifas.ufl.edu/PI052>.
- Muhammad, W., Iqbal, N., Saeed, S. & Javed, M. 2013. Monitoring and varietal preference of mango midge, *Procontarinia mangicola* (Diptera: Cecidomyiidae). *Pakistan Journal of Zoology* 45(5): 1273-1278.
- Muhammad, W., Javed, M., Saeed, S., Kassi, A.K., Iqbal, N. & Ahmad, I., 2017. Evaluation of different insecticides against mango midges (Diptera: Cecidomyiidae). *Journal of Entomology and Zoology Studies* 5(4): 1888-1890.
- Muller, H.R.A., 1940. Overzicht van de belangrijkste mangga-ziekten in Nederlandsch Indie. (English abstract in *Review of Applied Mycology*, 19: 355.
- Nachiappan, R.M. & Baskaran, P., 1986. Field evaluation of certain insecticidal sprays against mango leaf-hoppers. *Pesticides* 20: 41–44.
- Nafus, D., 1991. Biological control of *Penicillaria jocosatrix* (Lepidoptera: Noctuidae) on mango on Guam with notes on the biology of its parasitoids. *Environmental Entomology* 20: 1725–1731.
- Narashimhan, U. A. & Chacko, M. J., 1988. The distribution of some *Rastrococcus* spp. (Homoptera: Pseudococcidae) on mango in India. *Bulletin of Entomology Research* 81: 445–448.
- Nascimento, A., Malavasi, A., Morgante, J. and Duarte, A.L., 1992. Hot-water immersion treatment for mango infested with *Anastrepha fraterculus*, *A. obliqua*, and *Ceratitidis capitata* (Diptera: Tephritidae) in Brazil. *Journal of Economic Entomology* 85: 456–460.
- Nasir, M., Muhammad Mughal, S., Mukhtar, T. & Zaman Awan, M., 2014. Powdery mildew of mango: A review of ecology, biology, epidemiology and management. *Crop Protection* 64: 19–26.

- Navarro-Llopis, V., Primo, J. & Vacas, S., 2012. Efficacy of attract-and-kill devices for the control of *Ceratitis capitata*. *Pest Manag. Sci.* 69: 478–482.
- Ndlela, S., Mohamed, S., Ndegwa, P.N., Amo, G.O. & Ekesi, S., 2016. Male annihilation technique using methyl eugenol for field suppression of *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae) on mango in Kenya. *Afr. Entomol.* 24: 437–447.
- Neergaard, P., 1945. Danish species of *Alternaria* and *Stemphylium*. Taxonomy, Parasiticism, Economical Significance. Einar Munksgaard, Copenhagen, Denmark.
- Nofal, M.A. & Haggag, W.A., 2006. Integrated management of powdery mildew of mango in Egypt. *Crop Protection* 25: 480-486.
- Nunez-Elisea, R., 1985. Flowering and fruit set of a monembryonic and a polyembryonic mango as influenced by potassium nitrate sprays and shoot decapitation. *Proceedings of the Florida State Horticultural Society* 98: 179–183.
- Nunez-Elisea, R., Davenport, T.L. & Caldeira, M.L., 1993. Bud initiation and morphogenesis in ‘Tommy Atkins’ mango as affected by temperature and triazole growth retardants. *Acta Horticulturae* 341: 192–198.
- Obra, G.B., Resilva, S.S., Follet, P.A. & Lorenzana, L.R.J., 2014. Large-Scale Confirmatory Tests of a Phytosanitary Irradiation Treatment Against *Sternochetus frigidus* (Coleoptera: Curculionidae) in Philippine Mango. *Journal of Economic Entomology* 107(1): 161-165.
- Okigbo, R.N., 2001. Occurrence, pathogenicity and survival of *Macrophoma mangiferae* in leaves, branches and stems of mango (*Mangifera indica* L.). *Plant Protection Science* 37: 138–144.
- Okigbo, R.N. & Osuinde, M.I., 2003. Fungal leaf spot diseases of mango (*Mangifera indica* L.) in southeastern Nigeria and biological control with *Bacillus subtilis*. *Plant Protection Science* 39: 70–77.
- Otero-Colina, G., Rodriguez-Akvarado, G., Fernandez-Pavia, S., Maymon, M., Ploetz, R.C., Aoki, T., O’Donell, K. & Freeman, S., 2010. Identification and characterization of a novel etiological agent of mango malformation disease in Mexico, *Fusarium mexicanum* sp. nov. *Phytopathology* 100: 1176-1184.
- Ovruski, S., Aluja, M., Sivinski, J. & Wharton, R., 2000. Hymenopteran parasitoids on fruit infesting Tephritidae (Diptera) in Latin America and the southern United States: diversity, distribution, taxonomic status and their use in fruit fly biological control. *Integrated Pest Management Reviews* 5: 81–107.
- Palo, M.A., 1933. Sclerotium seed rot and seedling rot of mango. *Philippine Journal of Science* 52: 237–261.
- Palti, J., Pinkas, Y. & Chorin, M., 1974. Powdery mildew of mango. *Plant Disease Report* 58: 45–49.
- Park, H. and M. Duponte. 2008. How to cultivate indigenous microorganisms. *CTAHR Cooperative Extension Service Publication*. BIO-9. 7 pp.
- Patino-Vera, M., Jimenez, B., Balderas, K., Ortiz, M., Allende, R., Carrillo, A. & Galindo, E., 2005. Pilot-scale production and liquid formulation of *Rhodotorula minuta*, a potential biocontrol agent of mango anthracnose. *Journal of Applied Microbiology* 99: 540–50.
- Peak, E.M., Fitzell, R.D., Hannah, R.S. & Batten, D.J., 1986. Development of a microprocessor-based data recording system for predicting plant disease based on studies on mango anthracnose. *Computers and Electronics in Agriculture* 1: 251–262.

- Peck, S.L. & McQuate, G.T., 2000. Field tests of environmentally friendly malathion replacements to suppress wild Mediterranean fruit fly (Diptera: Tephritidae) populations. *Journal of Economic Entomology* 93: 280–289.
- Pedigo, L. & Buntin, D. (editors), 1993. "Handbook of Sampling Methods for Arthropods in Agriculture". Preface. CRC Press, Boca Raton, Florida.
- Pelikan, W. & Unger, G., 1971. The activity of potentized substances. *British Homoeopathy Journal*, 60: 232-266.
- Peng, R. & Christian, K., 2005. The control efficacy of the weaver ant, *Oecophylla smaragdina* (Hymenoptera: Formicidae) on the mango leafhopper, *Idioscopus nitidulus* (Hemiptera: Cicadellidae) in mango orchards in the Northern Territory. *International Journal of Pest Management*, 51: 297–304.
- Peng, R.K. & Christian, K., 2006a. Effective control of Jarvis's fruit fly *Bactrocera jarvisi* (Diptera: Tephritidae), by weaver ant, *Oecophylla smaragdina* (Hymenoptera: Formicidae), in mango orchards in the Northern Territory of Australia. *International Journal of Pest Management* 52: 275–282.
- Peng, R. K. & Christian, K. 2006b. Potential for organic mango production in the north territory of Australia using weaver ant, *Oecophylla smaragdina* (Hymenoptera: Formicidae). *Acta Horticulturae* 767: 81–88.
- Peña J.E., 1993. Pests of mango in Florida. *Acta Hort* 341: 395–406.
- Peña, J.E., 2002. Integrated pest management and monitoring techniques for mango pests. VII Int Mango Symp 645: 151–161.
- Peña, J. E., 2004. Integrated pest management and monitoring techniques for mango pests. *Acta Horticulturae* 645: 151–61.
- Peña, J.E. & Mohyuddin, A.I., 1997. Insect Pests. In: "The Mango. Botany, Production and Uses", edited by Litz, R.E. Published by Cab International, UK. Pp. 327-362.
- Peña J.E., Mohyuddin A.I. & Wysoki, M., 1998. A review of the pest management situation in mango agroecosystems. *Phytoparasitica* 26: 129–148.
- Peña, J.E., Duncan, R., Vasquez, T. & Hennessey, M., 1999. Guava arthropod seasonality and control of fruit flies in south Florida. *Proceedings of the Florida State Horticultural Society* 112: 206–209.
- Peña, J.E., Aluja, M. & Wysoki, M., 2009. Pests. In: "The Mango. Botany, Production and Uses. 2nd Edition.", edited by Litz, R.E. Published by Cab International, UK. Pp. 317-366.
- Pinero, J., Aluja, M., Vazquez, A., Equihua, M. & Varon, J., 2003. Human urine and chicken feces as fruit fly (Diptera: Tephritidae) attractants for resource-poor fruit growers. *Journal of Economic Entomology* 96: 334–340.
- Pinto, A.C.Q., Andrade, S.R.M., Amaro, A.A. & Gomes, U., 2004. Mango industry in Brazil. *Acta Horticulturae* 645: 37–50.
- Pinto, L., Ippolito, A. & Baruzzi, F., 2015. Control of spoiler *Pseudomonas* spp. on fresh cut vegetables by neutral electrolyzed water. *Food Microbiology* 50: 102–108.
- Plan, M.R.R., Joyce, D.C., Ogle, H.J. & Johnson, G.I., 2002. Mango stem-end rot (*Botryosphaeria dothidea*) disease control by partial-pressure infiltration of fungicides. *Australian Journal of Experimental Agriculture* 42: 625–629.
- Ploetz, R.C., 2001. Malformation: a unique and important disease of mango, *Mangifera indica* L. In: "Fusarium: Paul E. Nelson Memorial Symposium", edited by Summerell, B.A., Leslie, J.F., Backhouse, D.

and Bryden, W.L. Published by American Phytopathological Society (APS) Press, St Paul, Minnesota, USA. Pp. 233–247.

Ploetz, R. C., 2003. Diseases of mango. In: “Diseases of Tropical Fruit Crops”, edited by R.C. Ploetz. Published by CABI Publishing, Wallingford, UK. Pp. 327-363.

Ploetz, R.C., 2004. The major diseases of mango: strategies and potential for sustainable management. *Acta Horticulturae* 645: 137–150.

Ploetz, R.C., 2017. The mango diseases trilogy. Proc. XI International Mango Symposium. *Acta Horticulturae* 1183: 221-227.

Ploetz, R.C., 2018. Integrated disease management in mango cultivation. In: “Achieving Sustainable Cultivation of Mangoes”, edited by V. Galán-Saúco & P. Lu. Burleigh Dodds Series in Agriculture Science, Number 34. Published by Burleigh Dodds Science Publishing Limited, Cambridge, UK. Pp. 459-496.

Ploetz, R.C. & Freeman, S., 2009. Foliar, floral and soilborne diseases. In: “The Mango: Botany, Production and Uses. 2nd Edition”, edited by R. Litz. Published by CAB International, Oxfordshire, UK. Pp. 231-302.

Ploetz, R.C. & Prakash, O., 1997. Foliar, floral and soilborne diseases of mango. In: “The Mango: Botany, Production and Uses”, edited by R. Litz. Published by CAB International, Wallingford, UK. Pp. 281–325.

Ploetz, R.C., Benschler, D., Vasquez, A., Colls, A., Nagel, J. & Schaffer, B., 1996a. A re-evaluation of mango decline in Florida. *Plant Disease* 80: 664–668.

Ploetz, R. C, Vazquez, A. & Benschler, D., 1996b. First report of *Fusarium decemcellulare* as a pathogen of mango in the United States. *Plant Disease* 80: 1207.

Poinar, G.O., Jr & Hislop, R.G., 1981. Mortality of Mediterranean fruit fly adults (*Ceratitis capitata*) from parasitic nematodes (*Neoaplectana* and *Heterorhabditis* spp.). *Institute for Research and Cognitive Science, Medical Science. Microbiology, Parasitology and Infectious Disease* 9: 641.

Prakash, O., 2004. Diseases and disorders of mango and their management. In: “Diseases of Fruits and Vegetables. Volume I”, edited by S.A.M.H. Naqvi. Published by Kluwer Academic Publishers, The Netherlands. Pp. 511-619.

Prakash, O., 2012. IPM schedule for mango pests. National Horticulture Mission. Ministry of Agriculture. Department of Agriculture and Cooperation. Krishi Bhawan, New Delhi.

Prakash, O. & Misra, A.K., 1986. Evaluation of Mango Germplasm Against Powdery Mildew Under Natural Condition. Annual Report. CIHNP, Lucknow. Pp. 62-63.

Prakash, O. & Misra, A. K., 1992. Important diseases of mango and their effect on production. *Biological Memoirs* 18: 39–55.

Prakash, O. & Misra, A. K., 1993a. Fungal diseases of sub-tropical horticultural fruit crops. In: “Advances in Horticulture”, edited by K.L. Chadha. Published by Malhotra Publishing House, New Delhi (India). Pp. 1275–372.

Prakash, O. & Misra, A. K., 1993b. Integrated approach in the management of mango diseases (Abstract). In: “National Conference on Eco-friendly Approaches in the Management of Pests/ Diseases and Industrial Effluents”. 20–22 December at CSAU & T, Kanpur. Pp. 70–71.

Prakash & Patil, R.R., 2018a. Field evaluation of new insecticides against scale insect, *Hemilecanium imbricans*, (Coccidae: Homoptera) on mango. *Journal of Entomology and Zoology Studies* 6(2): 2615-2620.

- Prakash & Patil, R.R., 2018b. Evaluation of different insecticides and biorationals against scale insect, *Hemilecanium imbricans* (Green), under laboratory conditions. *Int J Curr Microbiol App Sci* 7(7): 866-881.
- Prakash, O. & Raoof, M.A., 1994. Studies on powdery mildew (*Oidium mangiferae*) disease of mango: distribution, perpetuation, losses and chemical control. *Biological Memoirs* 20: 31-45.
- Prakash, O and Singh, U.N., 1976. New Disease of mango. *Proceedings Fruit Research Workshop, Hyderabad*, May, 24-28th. Pp. 300-302.
- Prakash, O. & Singh, U.N., 1977. Phoma blight, a new disease of mango (*Mangifera indica* L.). *Plant Disease Report* 61: 419-421.
- Prakash, O. & Srivastava, K.C., 1987. Mango Diseases and their Management- A World Review. Tommorrow's Printer, New Delhi, India.
- Prusky, D., 1988. The use of antioxidants to delay the onset of anthracnose and stem end rot in avocado fruits after harvest. *Plant Disease* 72: 381–384.
- Prusky, D., 1996. Pathogen quiescence in postharvest diseases. *Annual Reviews of Phytopathology* 34: 413-434.
- Prusky, D. & Keen, N.T., 1993. Involvement of preformed antifungal compounds in the resistance of subtropical fruits to fungal decay. *Plant Disease* 77: 114–119.
- Prusky, D. & Lichter, A., 2007. Activation of quiescent infections by postharvest pathogens during transition from the biotrophic to the necrotrophic stage. *FEMS Microbiological Letters* 268: 1–8.
- Prusky, D. & Plumbley, R.A., 1992. Quiescent infections of *Colletotrichum* in tropical and subtropical fruits. In: "Colletotrichum: Biology, Pathology and Control", edited by Bailey, J.A. and Jeger, M.J. Published by CAB International, Wallingford, UK. Pp. 289–307.
- Prusky, D. & Yakoby, N., 2003. Pathogenic fungi: leading or led by ambient pH?. *Molecular Plant Pathology* 4: 509–516.
- Prusky, D., Fuch, Y. & Yanko, U., 1983. Assessment of latent infections as a basis for control of post harvest disease of mango. *Plant Disease* 67: 816–818.
- Prusky, D., Plumbley, R. & Kobiler, I., 1991. Modulation of natural resistance of avocado fruits to *Colletotrichum gloeosporioides* by CO₂. *Physiological and Molecular Plant Pathology* 39: 325–334.
- Prusky, D., Gat, Z. & Burd, P., 1992. Effect of relative humidity during mango growth on the incidence of quiescent infections of *Alternaria alternata*. *Plant Disease* 77: 249–252.
- Prusky, D., Freeman, S., Rodriguez, R.J. & Keen, N.T., 1993. A nonpathogenic mutant strain of *Colletotrichum magna* induces resistance to *C. gloeosporioides* in avocado fruits. *Molecular Plant-Microbe Interactions* 7: 326–333.
- Prusky, D., Falik, E., Kobiler, I., Fuchs, Y., Zauberman, G., Pesis, E., Roth, I., Weksler, A., Akerman, M. & Ykutiely, O., 1996. Hot water brush: a new method for the control of post-harvest disease caused by alternaria rot in mango fruits. *Acta Horticulturae* 455: 780–785.
- Prusky, D., Fuchs, Y., Kobiler, I., Roth, I., Weksler, A., Shalom, Y., Fallik, E., Zauberman, G., Pesis, E., Akerman, M., Ykutiely, O., Weisblum, A., Regev, R. & Artes, L., 1999. Effect of hot water brushing, prochloraz treatment and waxing on the incidence of black spot decay caused by *Alternaria alternata* in mango fruit. *Postharvest Biology and Technology*, vol. 15: 165–174.

- Prusky, D., McEvoy, J.L., Leverentz, B. & Conway, L.S., 2001. Local modulation of host pH by *Colletotrichum* species as a mechanism to increase virulence. *Molecular Plant Microbe Interactions* 14: 1105–1113.
- Prusky, D., Shalom, Y., Kobiler, I., Akerman, M. & Fuchs, Y., 2002. The level of quiescent infection of *Alternaria alternata* in mango fruits at harvest determines the postharvest treatment applied for the control of rots during storage. *Postharvest Biology and Technology* 25: 339–347.
- Prusky, D., Kobiler, I., Akerman, M. & Miyara, I., 2006. Effect of acidic solutions and acidic prochloraz on the control of postharvest decay caused by *Alternaria alternata* in mango and persimmon fruits. *Postharvest Biology and Technology* 42: 134–141.
- Prusky, D., Kobiler, I., Miyara, I. & Alkan, N., 2009. Fruit diseases. In: “The Mango: Botany, Production and Uses. 2nd Edition”, edited by R. Litz. Published by CAB International, Oxfordshire, UK. Pp. 210-230.
- Pruvost, O., Couteau, A. & Luisetti, J., 1989. Efficacite de differentes formulations chimiques pour Jutter contre la maladie des taches noires de la mangue (*Xanthomonas campestris* pv. *mangiferaeindicae*). *Fruits* 44: 343–50.
- Ramaiah, A.K. & Raj Kumar, H., 2015. In Vitro Antifungal activity of some plant extracts against *Fusarium oxysporum* f. sp. *Lycopersici*. *Asian Journal of Plant Science and Research* 5(1): 22-27.
- Ramos, L.J., Lara, S.P., McMillan, R.T. & Narayanan, K.R., 1991. Tip dieback of mango (*Mangifera indica*) caused by *Botryosphaeria ribis*. *Plant Disease* 75, 315–318.
- Rao, B.S., 1975. *Maladies of Hevea in Malaysia*. Rubber Research Institute of Malaysia, Kuala Lumpur, Malaysia.
- Reckhaus, P. & Adamou, I., 1987. Hendersonula dieback of mango in Niger. (Disease note). *Plant Disease* 71: 1045.
- Reddy, G.S., Govinda, R.P. & Paparao, A., 1961. Black banded disease of mango. *Andhra Agricultural Journal* 8: 120–123.
- Reddy, P.V.R., Chakravarthy, A.K., Sudhagar, S. & Kurian, R., 2014. A simple technique to capture, contain and monitor the fresh-emerging beetles of tree borers. *Curr Biotica* 8(2): 191–194.
- Reddy, P.V.R., Gundappa, P. & Chakravarthy, A.K., 2018. Pests of mango. In: “Pests and Their Management”, edited by Omkar. Published by Springer Nature Signature. Pp. 415-440. https://doi.org/10.1007/978-981-10-8687-8_12.
- Reuveni, M., 2000. Efficacy of trifloxystrobin (Flint), a new strobilurin fungicide, in controlling powdery mildew on apple, mango and nectarine and rust on prune trees. *Crop Protection* 19: 335-341.
- Rey, L., 2003. Thermoluminescence of ultra-high dilutions of lithium chloride and sodium chloride. *Physica A: Statistical Mechanics and its Applications*, 323: 67–74.
- Ribiero, I.J.A., 1980. Seca de mangueira. Agentes causais e estudio da molesta. *Anais do I Simposio Brasileiro Sobre a Cultura de Mangueira*. Sociedade Brasileira de Fruticultura, Jaboticobal, Brazil. Pp. 123–130.
- Ribiero, I.J.A., 1997. Doencas da mangueira (*Mangifera indica* L.). In: “Manual de Fitopatologia, Volume 2: Doencas das Plantas Cultivadas”, edited by Kimati, H., Amorim, A., Bergamin-Filho, A., Camargo, L. E. A. and Rezende, J. A. M. Published by Agronomica Ceres, Sao Paulo, Brazil.
- Ribiero, I. J. A., Rosetto, C. J., Donadio, L. C., Sabino, J. C. Martins, A. L. M. & Gallo, P. B., 1995. Mango wilt. XIV Selection of mango (*Mangifera indica* L.) rootstocks resistant to the mango wilt fungus *Ceratocystis fimbriata* Ell. & Halst. *Acta Horticulturae* 370: 159–161.

- Robacker, D.C., 2001. Roles of putrescine and 1-pyrroline in attractiveness of technical-grade putrescine to the Mexican fruit fly (Diptera: Tephritidae). *Florida Entomologist* 84: 679–685.
- Robacker, D.C. & Warfield, W.C., 1993. Attraction of both sexes of Mexican fruit fly, *Anastrepha ludens*, to a mixture of ammonia, methylamine, and putrescine. *Journal of Chemical Ecology* 19: 2999–3016.
- Robacker, D.C., Martinez, A.J., Garcia, J.A., Diaz, M. & Romero, C., 1996. Toxicity of *Bacillus thuringiensis* to Mexican fruit fly (Diptera: Tephritidae). *Journal of Economic Entomology* 89: 104–110.
- Robacker, D.C., DeMilo, A.B. & Voaden, D.J., 1997. Mexican fruit fly attractants: effects of 1-pyrroline and other amines on attractiveness of a mixture of ammonia, methylamine, and putrescine. *Journal of Chemical Ecology* 23: 1263–1280.
- Rossetto, C. J., Ribeiro, I. J. A., Igue, T. & Gallo, P. B., 1996. Mango wilt: XV. Varietal resistance against two isolates of *Ceratocystis fimbriata*. *Bragantia* 55: 117–121.
- Rossi, F., Azevedo-Filho, J.A., Melo, P.C.T., Ambrosiano, E.J., Guirado, N. & Schammas, E.A., 2007. Cultivo orgânico de batata com aplicação de preparados homeopáticos. In: Resumos del 5º Congresso Brasileiro de Agroecologia. *Rev. Bras. de Agroecologia*, 2: 937-940.
- Ruehle, G.D. & Ledin, R.B., 1955. *Mango Growing in Florida*. Bulletin 574. Agricultural Experiment Station, University of Florida, Gainesville, Florida.
- Rungjindamai, N., 2016. Isolation and evaluation of biocontrol agents in controlling anthracnose disease of mango in Thailand. *Journal of Plant Protection Research* 56 (3): 306-311.
- Sadana, G.L. & Kumari, M., 1991. Predatory potential of the lysomanid spider, *Lyssomanes sikkimeri* Tikader on the mango hopper, *Ideoscopus clipealis* (Lethierry). *Entomon* 16: 283–285.
- Sangchote, S., 1998. Fruit rots of mangosteen and their control. In: “Post-harvest Handling of Tropical Fruit”, edited by Champ, B.R., Highley, E. and Johnson, G.I. Australian Centre for International Agricultural Research (ACIAR), Canberra. Pp. 81–86.
- Santos, A. M. G., Lins, S. R. O., da Silva, J. M. & de Oliveira, S. M. A., 2015. Low doses of gamma radiation in the management of postharvest *Lasiodiplodia theobromae* in mangos. *Brazilian Journal of Microbiology* 46: 841–847.
- Schaffer, B., Larson, K.D., Snyder, G.H. & Sanchez, C.A., 1988. Identification of mineral deficiencies associated with mango decline by DRIS. *HortScience* 23: 617–619.
- Schirra, M., D'Hallewin, G., Ben-Yehoshua, S. & Fallik, E., 2000. Host–pathogen interactions modulated by heat treatment. *Postharvest Biology and Technology* 21: 71–85.
- Schoeman, M.H., Zulu, N.B., Botha, F.A. & Calitz, F.J. 2018. Integrated control of mango blossom malformation in South Africa. *South African Journal of Plant and Soil*: 36, 51-56.
- Schreiner, I., 1987. Mango shoot caterpillar control on mango flowers, 1985. *Insecticide and Acaricide Tests* 12: 94.
- Secretariat Commonwealth, 1987. The development and adaptation of methods for control of anthracnose in mango: 29-38.
- Senghor, A. L., Sharma, K., Kumar, P. L. & Bandyopadhyay, R., 2012. First report of mango malformation disease caused by *Fusarium tuiense* in Senegal. *Plant Disease* 96: 1582.

- Shah, A.H., Jhala, R.C., Patel, G.M. & Patel, S.H., 1983. Evaluation of effective dose of monocrotophos for the control of mango hopper *Amritodus atkinsoni* Lethierry (Cicadellidae: Homoptera) by injection method and its comparison with foliar application. *Gujarat Agricultural University Research Journal* 9: 14–18.
- Sharma, G., Kumar, N., Weir, B.S., Hyde, K.D., & Shenoy, B.D., 2013. The ApMat marker can resolve *Colletotrichum* species: a case study with *Mangifera indica*. *Fungal Diversity* 61 (1): 117–138
- Sharp, J.L., Ouye, M.T., Thalman, R., Hart, W.G., Ingle, S.J. & Chew, V., 1988. Submersion of 'Francis' mango in hot water as a quarantine treatment for the West Indian fruit fly and the Caribbean fruit fly (Diptera: Tephritidae). *Journal of Economic Entomology* 81: 1431–1436.
- Sharp, J.L., Ouye, M.T., Ingle, S.J., Hart, W.G., Enkerlin, E., Celedonio, H., Toledo, J., Stevens, L., Quintero, E., Reyes, J. & Schwarz, A., 1989a. Hot-water quarantine treatment for mangoes from the State of Chiapas, Mexico, infested with Mediterranean fruit fly and *Anastrepha serpentina* (Wiedemann) (Diptera: Tephritidae). *Journal of Economic Entomology* 82: 1663–1666.
- Sharp, J.L., Ouye, M.T., Hart, W.G., Ingle, S.J., Hallman, G., Gould, W. & Chew, V., 1989b. Immersion of Florida mangoes in hot water as a quarantine treatment for Caribbean fruit fly (Diptera: Tephritidae). *Journal of Economic Entomology* 82: 186–188.
- Sharp, J.L., Ouye, M.T., Ingle, S.J. & Hart, W.G., 1989c. Hot water quarantine treatment for mangoes from Mexico infested with Mexican fruit fly and West Indian fruit fly (Diptera: Tephritidae). *Journal of Economic Entomology* 82: 1657–1662.
- Shellie, K.C. & Mangan, R.L., 2002a. Hot water immersion as a quarantine treatment for large mangoes: artificial versus cage infestation. *Journal of the American Society for Horticultural Science* 127: 430–434.
- Shellie, K.C. & Mangan, R.L., 2002b. Cooling method and fruit weight: efficacy of hot water quarantine treatment for Mexican fruit fly (Diptera: Tephritidae) in mango. *HortScience* 37: 910–913.
- Shivananda, T.N., Verghese, A., Hegde, R.M., Kumar, N.K.K. & Singh, A.S., 2012. Management of mango trunk borers using sealer cum healer. Technical bulletin no. 72/12, IIHR, Bangalore.
- Shukla, R.P. & Prasad, V.G., 1984. Evaluation of some insecticides against mango hopper, *Idioscopus clypealis* (Lethierry) (Homoptera: Cicadellidae). *Indian Journal of Agricultural Science* 54: 677–681.
- Shukla, R.P. & Tandon, P.L., 1985. Bio-ecology and management of the mango weevil, *Sternochetus mangiferae* (Fabricius) (Coleoptera: Curculionidae). *International Journal of Tropical Agriculture* 3: 293–303.
- Shukla, R. P., Tandon, P. L. & Suman, C. L., 1988. Intra-tree distribution of eggs and diapausing adults of the stone weevil. *Acta Horticulturae* 231: 566–70.
- Silimela, M. & Korsten, L., 2007. Evaluation of pre-harvest *Bacillus licheniformis* sprays to control mango fruit diseases. *Crop Protection* 26: 1474–81.
- Silveira, S. F., Harrington, T. C., Mussi-Dias, V., Engelbrecht, C. J. B., Alfenas, A. C. & Silva, C. R., 2006. *Annona squamosa*, a new host of *Ceratocystis fimbriata*. *Fitopatologia Brasileira* 31: 394–397.
- Singh, G., 1991. Loss assessment, ecology and management of mango fruit fly, *Dacus* sp. *Acta Horticulturae* 291: 425–436.
- Singhania, P.K. & Singhania, A., 2014. Homoeopathy in agriculture. In Ahmann, G. & Aksoy, U. (eds.). Proceedings of the 4th ISOFAR Scientific Conference. 'Building Organic Bridges', at the Organic World Congress, Istanbul, Turkey Pp. 667-670.

- Sinniah, G. D., Adikaram, N. K. B., Vithanage, I. S. K., Abayasekara, C. L., Maymon, M. & Freeman, S., 2013. First report of mango malformation disease caused by *Fusarium mangiferae* in Sri Lanka. *Plant Disease* 97: 427.
- Sivankalyani, V., Feygenberg, O., Diskin, S., Wright, B. & Alkanm, N., 2016. Increased anthocyanin and flavonoids in mango fruit peel are associated with cold and pathogen resistance. *Postharvest Biology and Technology*, vol. 111: 132–139.
- Sivinski, J.M., 1996. The past and potential of biological control of fruit flies. In: "Fruit Fly Pests. A World Assessment of their Biology and Management", edited by McPherson, B.A. and Steck, G.J. Published by St Lucie Press, Delray Beach, Florida. Pp. 365–375.
- Sivinski, J.M., Calkins, C.O., Baranowski, R.M., Harris, D., Brambila, J., Diaz, J., Bums, R.E., Holler, T. & Dodson, D., 1996. Suppression of Caribbean fruit fly (*Anastrepha suspensa* (Loew) Diptera: Tephritidae) population through releases of the parasitoid *Diachasmimorpha longicaudata* (Ashmead) (Hymenoptera: Braconidae). *Biological Control* 6: 177–185.
- Sivinski, J., Aluja, M. & Lopez, M., 1997. Spatial and temporal distributions of parasitoids of Mexican *Anastrepha* species (Diptera: Tephritidae) within the canopies of fruit trees. *Annals of the Entomological Society of America* 90: 604–618.
- Sivinski, J., Pinero, J. & Aluja, M., 2000. The distributions of parasitoids (Hymenoptera) of *Anastrepha* fruit flies (Diptera: Tephritidae) along an altitudinal gradient in Veracruz, Mexico. *Biological Control* 18: 258–269.
- Smith, K. 2012. Homeopathy is unscientific and unethical". *Bioethics* 26 (9): 508–512.
- Snowdon, A.L., 1990. Color Atlas of Post-Harvest Diseases and Disorders of Fruits and Vegetables. Vol. 1. General Introduction and Fruits. CRC Press, Boca Raton, Florida.
- Soto-Manitiu, J., Jiron, L. & Hernandez, R., 1987. Chemical control and ecological observations of fruit flies of the genus *Anastrepha* Schiner (Diptera: Tephritidae) on mango. *Turrialba* 37: 245–251.
- Srivastava, R.P., 1998. Pests of mango. In: "Mango Cultivation", edited by Srivastava, R.P. International Book Distributing, Charbagh, Lucknow, India. Pp. 175–299.
- Srivastava, R. & Tandon, P., 1986. Natural occurrence of two entomogenous fungi pathogenic to mango hopper, *Idioscopus clypealis* Leth. *Indian Journal of Plant Pathology* 4: 121–123.
- Stark, J.D., Vargas, R. & Miller, N., 2004. Toxicity of spinosad in protein bait to three economically important tephritid fruit fly species (Diptera: Tephritidae) and their parasitoids (Hymenoptera: Braconidae). *Journal of Economic Entomology* 97: 911–915.
- Steck, G.J., 2003. Mango Fruit Fly, Marula Fruit Fly, *Ceratitis cosyra* (Walker) (Insecta: Diptera: Tephritidae). Available at: <http://edis.ifas.ufl.edu/IN563> (accessed 20 July 2007).
- Steiner, L.F. & Lee, R.K., 1955. Large area tests of male annihilation method for oriental fruit fly control. *Journal of Economic Entomology* 48: 311–317.
- Steiner, L.F., Mitchell, W.C., Harris, E., Kozuma, T. & Fujimoto, M., 1965. Oriental fruit fly eradication by male annihilation. *Journal of Economic Entomology* 58: 961–964.
- Summanwar, A. S, Raychaudhuri, S. P. & Phatak, S. C., 1966. Association of the fungus *Fusarium moniliforme* Sheld. with the malformation in mango (*Mangifera indica* L.). *Indian Phytopathology* 19: 227–228.

- Sutantawong, M., Orankanok, W., Enkerlin, W.R., Wornoaayporn, V. & Caceres, C., 2002. The sterile insect technique for control of the oriental fruit fly, *Bactrocera dorsalis* (Hendel), in mango orchards in Ratchaburi Province, Thailand. *Proceedings of 6th International Fruit Fly Symposium*, Stellenbosch, South Africa. Pp. 223–232
- Swirski, E., Wysoki, M. & Izhar, Y., 2002. Subtropical pests of Israel. Fruit Board of Israel and Institute of Plant Protection, The Volcani Center, Tel Aviv, 284 pp.
- Syed, R.A., Ghani, M.A. & Murtaza, M., 1970. Studies on trypetids and their natural enemies in West Pakistan. III. *Dacus (Strumeta) zonatus* (Saunders). *Technical Bulletin of the Commonwealth Institute of Biological Control* 13: 1–16.
- Syed Ab Rahman, S. F., Singh, E., Pieterse, C. M. J., & Schenk, P. M., 2018. Emerging microbial biocontrol strategies for plant pathogens. *Plant Sci.* 267: 102–111. doi: 10.1016/j.plantsci.2017.11.012.
- Sztejnberg, A., Galper, S., Mazar, Shlomit & Lisker, N., 1989. *Ampelomyces quisqualis* for biological and integrated control of powdery mildews in Israel. *Journal of Phytopathology* 124: 285–295.
- Tandon, P.L. & Lai, B., 1979. Studies on the chemical control of the mango hopper, *Idioscopus clypealis* Leth. (Cicadellidae: Homoptera). *International Pest Control* 21: 6–9.
- Tandon P.L. & Verghese, A., 1985. World list of insect, mite and other pests of mango, Technical document, vol 5. Indian Institute of Horticultural Research, Bangalore.
- Tarnowski, T., 2009. Using molecular analysis to investigate phylogenetic relationships in two tropical pathosystems: witches' broom of cacao, caused by *Moniliophthora perniciosa*, and mango anthracnose, caused by *Colletotrichum* spp. PhD dissertation (Gainesville, FL, USA: University of Florida).
- Tefera, A., Seyoum, T. & Woldetsadik, K., 2007. Effect of disinfection, packaging, and storage environment on the shelf life of mango. *Biosystems Engineering* 96: 201–212.
- Tenorio, A.M., Lizarondo, R.C. & Inocencio, T.Q., 1989. Integrated pest management (IPM) approach for the control of mango twig borer, *Niphonoclea capito* Pasc. Retrieved reference from <http://agris.fao.org/agris-search/search.do?recordID=PH8912575>.
- Terao, D., Campos, J. S. D., Benato, E. A. & Hashimoto, J. M., 2015. Alternative strategy on control of postharvest diseases of mango (*Mangifera indica* L.) by use of low dose of ultraviolet-C irradiation. *Food Engineering Reviews* 7: 171–175.
- Tiwari, R. K. S., Ashok, S., Rajput, M. L. & Bisen, R. K., 2006. Relative susceptibility of mango varieties to powdery mildew caused by *Oidium mangiferae*. *Advances in Plant Science* 19: 181–183.
- Toledo, J., Liedo, P., Williams, T. & Ibarra, J., 1999. Toxicity of *Bacillus thuringiensis* β -exotoxin to three species of fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology* 92: 1052–1056.
- Toledo, M., Stangarlin, J., Bonato, C., 2011. Homeopathy for the control of plant pathogens. *Physiology* 19(20): 21.
- Tsao, P. H., Luzaran, P. B., de los Santos, A. B., Portales, L. A., Gochango, A. M. & Gruber, L. C., 1994. Phytophthora crown and foot rot of mango detected in Philippine nurseries. *Plant Disease* 78: 100.
- Udayanga, D., Manamgoda, D.S., Liu, X., Chukeatirote, E., & Hyde, K.D., 2013. What are the common anthracnose pathogens of tropical fruits? *Fungal Diversity* 61 (1), 165–179.
- Uddin, M. N., Shefat, S.H.T., Mansura, M. & Moon, N.J., 2018. Management of Anthracnose Disease of Mango Caused by *Colletotrichum gloeosporioides*: A Review". *Acta Scientific Agriculture* 2.10: 169-177.

Umeya, K. & Hirao, J., 1975. Attraction of the jackfruitfly, *Dacus umbrosus* F. (Diptera: Tephritidae) and lace wing, *Chrysopa* sp. (Neuroptera: Chrysopidae) to lure traps baited with methyl eugenol and cuelure in the Philippines. *Applied Entomology and Zoology* 10: 60–62.

Upadhyay, S.K., Chaudhary, B. & Sapkota, B., 2013. Integrated management of mango stem borer (*Batocera rufomaculata* Dejan) in Nepal. *Glob J Biol Agric Health Sci* 2(4): 132–135.

USDA, 2019. Treatment Manual. Retrieved from: https://www.aphis.usda.gov/import_export/plants/manuals/ports/downloads/treatment.pdf.

Usha, K., Singh, B., Praseetha, P. Deepa, N., Agarwal, D.K., Agarwal, R. & Nagaraja, A., 2009. Antifungal activity of *Datura stramonium*, *Calotropis gigantea* and *Azadirachta indica* against *Fusarium mangiferae* and floral malformation in mango. *European Journal of Plant Pathology* 124: 637–657.

Van Mele, P., 2008. A historical review of research on the weaver ant *Oecophylla* in biological control. *Agricultural and Forestry Entomology*, 10: 13–22.

Van Mele, P., Vayssieres, J.-F., van Tellingen, E. & Vrolijk, J., 2007. Effects of an African weaver ant, *Oecophylla longinoda*, in controlling mango fruit flies (Diptera: Tephritidae) in Benin. *Journal of Economic Entomology* 100: 695–701.

Vanniere, H., Didier, C., Rey, J.Y., Diallo, T.M., Keita, S. & Sangare, M., 2004. La mangue en Afrique de l'Ouest francophone: les systemes de production et les itineraires techniques. *Fruits* 59: 383–398.

Vargas, R.I., Souder, S.K., Mackey, B., Cook, P., Morse, J.G. & Stark, J.D., 2012. Field trials of solid triple lure (trimedlure, methyl eugenol, raspberry ketone, and ddvp) dispensers for detection and male annihilation of *Ceratitidis capitata*, *Bactrocera dorsalis*, and *Bactrocera cucurbitae* (Diptera: Tephritidae) in Hawaii. *J. Econ. Entomol.* 105: 1557–1565.

Veeresh, G.K., 1989. Pest problems in mango – world situation. *Acta Horticulturae* 231: 551–565.

Veldman WM, Regnier, T. & Augustyn W., 2017. Biocontrol of *Fusarium mangiferae* responsible for mango malformation using bacterial isolates. *Scientia Horticulturae*, 234: 186-195.

Verghese, A. 2000. Effect of imidacloprid, Lambda-cyhalothrin and Azadirachtin on the mango hopper, *Indioscopus niveosparsus* (Leth.) (Homoptera: Cicadellidae). *Acta Horticulturae* 509: 733–735.

Verghese, A. & Devi Thangam. S., 2011. Mango hoppers and their management. Extension folder no: 71-11, ATIC series: 31-11. Indian Institute of Horticultural Research, Bangalore.

Verghese, A., Nagaraju, D.K., Jayanthi, P.D.K., Vasudev, V. & Madhura, H.S., 2004. Is azadirachtin useful in the management of the mango stone weevil (*Sternonchetus mangiferae* Fabricius)? *Journal of Food Agriculture and Environment* 2: 213–216.

Verma, O.P. & Singh, R.D., 1970. Epidemiology of mango dieback caused by *Botyodiplodia theobromae* Pat. *Indian Journal of Agricultural Science* 40, 813–818.

Viegas, A. P., 1960. Mango blight. *Bragantia* 19:163–82.

Viljoen, N. M. & Kotze, J. M., 1972. Bacterial black spot of mango. *Citrus Grow. Sub-Trop. Fruit J.* June: 5–8.

Villalobos, J., Flores, S., Liedo, P. & Malo, E.A., 2017. Mass trapping is as effective as ground bait sprays for the control of *Anastrepha* (Diptera: Tephritidae) fruit flies in mango orchards. *Pest Manag. Sci.* 73: 2105–2110.

- Vyas, R.V., Patel, J.J., Godhani, P.H. & Yadav, D.N., 1993. Evaluation of green muscardine fungus (*Metarrhizium anisopliae* var. *anisopliae*) for control of mango hopper (*Amritodus atkinsoni*). *Indian Journal of Agricultural Sciences* 63: 602–603.
- Waddell, W.J., Crooks, N.H. & Carmichael, P.L., 2004. Correlation of tumours with DNA adducts from methyl eugenol and tamoxifen in rats. *Toxicological Sciences* 79: 38–40.
- Wang, X.G., Jarjees, E.A., McGraw, B.K., Bokonon-Ganta, A.H., Messing, R.H. & Johnson, M.W., 2005. Effects of spinosad-based fruit fly bait GF-120 on tephritid fruit fly and aphid parasitoids. *Biological Control* 35: 155–162.
- Wang, K-H., DuPonte, M. & Chang, M., 2012. Use of Korean Natural Farming for Vegetable Crop Production in Hawaii. *The Food Provider*, Dec 2012 - Febr 2013. Pp. 7.
- Waterhouse, D.F., 1998. *Biological control of insect pests: South-East Asian prospects*. Australian Centre for International Agricultural Research Monograph No. 51, Canberra, Australia, 548 pp.
- Weir, B. S., Johnston, P. R. and Damm, U., 2012. The *Colletotrichum gloeosporioides* species complex. *Studies in Mycology* 73: 115–180.
- Wharton, R.A., 1978. Classical biological control of fruit-infesting Tephritidae. In: “Fruit Flies. Their Biology, Natural Enemies and Control. Vol. B”, editd by Robinson, A.S. and Hooper, G. Published by Elsevier, Amsterdam, the Netherlands. Pp. 303–313.
- Whitwell, A.C., 1993. The pest/predator/parasitoid complex on mango inflorescences in Dominica. *Acta Horticulturae* 341: 421–432.
- Wysocki M., Ben-Dov Y., Swirski E. & Izhar, Y., 1993. The arthropod pests of mango in Israel. *Acta Hortic.* 341: 452–466.
- Yadav, S.M., Patil, R.K., Balai, L.P. & Niwas, R., 2013. Post-harvest diseases of horticultural crops and their management. *Popular Kheti*, 1: 20-25.
- Yazdani, S.S. & Mehto, D.N., 1980. A note on the control of mango hopper, *Amritodus atkinsoni* Leth. at Dholi. *Indian Journal of Entomology* 42: 275–276.
- Yee, W., 1987. The mango in Hawaii. *Cooperative Extension Service, University of Hawaii Circular* 388: 19–22.
- Zagzog, O.A., Gad, M.M. & Hafez, N.H., 2017. Effect of nano-chitosan on vegetative growth, fruiting and resistance of malformation of mango. *Trends in Horticultural Research*, 7: 11-18.
- Zea-Bonilla, T., Martin-Sanchez, P. M., Hermoso, J. M., Carmona, M. P., Segundo, E. & Perez-Jimenez, R. M., 2007. First report of *Phytophthora citricola* on *Mangifera indica* in Spain. *New Disease Reports. Plant Pathology* 56: 356.
- Zhan, R.-L., Yang, S.-J., Ho, H.-H., Liu, F., Zhao, Y.-L., Chang, J.-M. & He, Y.-B., 2010. Mango malformation disease in South China caused by *Fusarium proliferatum*. *Journal of Phytopathology* 158: 721–725.
- Zhan, R.-L., Yang, S.-J., Liu, F., Zhao, Y.-L., Chang, J.-M. & He, Y.-B., 2012. First report of *Fusarium mangiferae* causing mango malformation in China. *Plant Disease* 96: 762.
- Zucchi, R.A., 2000. Especies de *Anastrepha*, sinonimias, plantas hospedeiras e parasitoides. In: “Moscas-frutas de Importancia Economica no Brasil: Conhecimento Basico e Aplicado”, edited by Malavasi, A. and Zucchi, R.A. Published by Holos Editora, Ribeirao Preto, Brazil.

